



On a Factorisation of Free Monoids

M. P. Schutzenberger

Proceedings of the American Mathematical Society, Vol. 16, No. 1 (Feb., 1965), 21-24.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9939%28196502%2916%3A1%3C21%3AOAFOFM%3E2.0.CO%3B2-4>

Proceedings of the American Mathematical Society is currently published by American Mathematical Society.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

ON A FACTORISATION OF FREE MONOIDS

M. P. SCHÜTZENBERGER

A property is given which relates two results of Spitzer [6]; it also relates two results of Chen, Fox and Lyndon [1]; the same remark applies to work of Meyer-Wunderli [5] and M. Hall [3] and to its generalisation by Lazard [4]. These connections are indicated more fully below.

In what follows, F is the free monoid generated by a fixed set X and F^+ denotes the set of all words of positive length of F . If the words f and f' of F belong to a submonoid F' of F , the words ff' and $f'f$ are said to be F' -conjugate. We consider the following conditions, I, I' and II, on a family $\{Y_j: j \in J\}$ of subsets of F^+ indexed by a totally ordered set J .

(I) (resp. (I')). Each $f \in F^+$ has at most (resp. at least) one representation in the form $f = f_1 f_2 \cdots f_n$, $n > 0$, where each $f_i \in Y_j$, and $j_1 \geq j_2 \geq \cdots \geq j_n$.

(II) Each F -conjugate class C has nonempty intersection with the submonoid F_j generated by Y_j for exactly one $j \in J$; further, $C \cap F_j$ is an F_j -conjugate class.

PROPOSITION 1. Any two of the three conditions I, I' and II imply the third one.

PROOF. Let \mathfrak{A} be the large algebra of F over the real field R . If U is a subset of F , we write $U = \sum \{f: f \in U\} \in \mathfrak{A}$. Since $(1 - U)^{-1} = 1 + \sum \{U^m: m > 0\}$, it follows that $(1 - U)^{-1} = G$ iff G is a submonoid freely generated by U .

Let us assume first that I and I' are satisfied; it follows that each F_j , $j \in J$, is freely generated by Y_j and that $(1 - X)^{-1} = \prod \{(1 - Y_j)^{-1}: j \in J\}$ where the product is taken according to the given ordering of J . Further, $\text{Log}(1 - X)^{-1} = \sum \{m^{-1} X^m: m > 0\} = \sum \{(\lambda f)^{-1} f: f \in F^+\}$ and $\text{Log}(1 - Y_j)^{-1} = \sum \{(\lambda_j f)^{-1} f: f \in F^+ \cap F_j\}$, where λf (resp. $\lambda_j f$) denotes the length of the word f with respect to the free basis X (resp. Y_j).

For each F -conjugate class C , let π_C denote the linear map of \mathfrak{A} onto R that satisfies $\pi_C f = 1$ if $f \in C$ and $\pi_C f = 0$ if $f \in F \setminus C$. Since π_C is constant on conjugate classes, for all $f', f'' \in F$ we have $\pi_C(f' f'') = \pi_C(f'' f')$; it follows that if $\mathfrak{L} \subset \mathfrak{A}$ is the large Lie algebra over R generated by F , then $\pi_C \mathfrak{L}' = 0$ for $\mathfrak{L}' = [\mathfrak{L}, \mathfrak{L}]$. According to our hy-

Received by the editors April 22, 1963 and, in revised form, June 13, 1963 and September 3, 1963.

pothesis, $\text{Log}(1 - \mathbf{X})^{-1} = \text{Log} \prod (1 - \mathbf{Y}_j)^{-1}$ whence, by the Campbell-Hausdorff formula $\text{Log}(1 - \mathbf{X})^{-1} = \sum \{ \text{Log}(1 - \mathbf{Y}_j)^{-1} : j \in J \} + K$ where $K \in \mathfrak{A}'$. Consequently,

$$(1) \quad \pi_C \text{Log}(1 - \mathbf{X})^{-1} = \sum \{ \pi_C \text{Log}(1 - \mathbf{Y}_j)^{-1} : j \in J \}.$$

If $f \in C$ has the form $f = g^p$ with maximal positive p , it follows that $p \text{ Card } C = \lambda C$ where λC is the common length of all $f \in C$; in particular, p is independent of the choice of $f \in C$. Now $\pi_C \text{Log}(1 - \mathbf{X})^{-1} = \sum \{ (\lambda f)^{-1} : f \in C \} = (\lambda C)^{-1} \text{Card } C = p^{-1}$. From (1) we conclude that

$$(2) \quad p^{-1} = \sum \{ (\lambda_j(C \cap F_j))^{-1} \text{Card } (C \cap F_j) : j \in J \}.$$

If $p = 1$, the sum in (2) can have only one nonzero term and II is verified for C . If $p > 1$, we conclude from the case $p = 1$ that g has an F -conjugate $g' \in F_{j_0}$ for some $j_0 \in J$. It follows that $f' = g'^p \in C \cap F_{j_0}$, that $C \cap F_{j_0}$ is an F_{j_0} -conjugate class and that $(\lambda_{j_0}(C \cap F_{j_0}))^{-1} \text{Card } (C \cap F_{j_0}) = p^{-1}$. It now follows from (2) that $C \cap F_j \neq \emptyset$ iff $j = j_0$, and the implication I & I' \Rightarrow II is verified.

Let us assume now that II is satisfied; it follows that for each $j \in J$ one has

$$(3) \quad \text{for any } f, f' \in F, \text{ if } ff', f'f \in F_j, \text{ then } f, f' \in F_j.$$

Consequently (cf., e.g., [2]), each F_j is freely generated by Y_j and (2), whence (1), holds for every F -conjugate class C . Let α be the natural homomorphism of \mathfrak{A} into the large algebra over R of the free commutative monoid generated by X . We deduce from (1) that $\alpha \text{Log}(1 - \mathbf{X})^{-1} = \sum \{ \alpha \text{Log}(1 - \mathbf{Y}_j)^{-1} : j \in J \}$, or, in equivalent fashion that $\alpha(1 - \mathbf{X})^{-1} = \alpha \prod \{ (1 - \mathbf{Y}_j)^{-1} : j \in J \}$. Now, I (resp. I') is equivalent to $S + \prod (1 - \mathbf{Y}_j)^{-1} = (1 - \mathbf{X})^{-1}$ where S (resp. $-S$) is an element of \mathfrak{A} in which every $f \in F$ has non-negative coefficient. Since $\alpha S = 0$ implies $S = 0$, the implication I & II \Rightarrow I' (resp. I' & II \Rightarrow I) is verified.

EXAMPLE 1. Let σ be a homomorphism of F into the additive group of R and identify J with R . For $r \in R$, let Y_r be the set of all $f \in F^+$ such that $\sigma f = r \lambda f$ and that $\sigma f' < r \lambda f'$ for every factorisation $f = f' f''$ ($f' \neq 1, f$). The fact that $\{ Y_r : r \in R \}$ satisfies I and I' (resp. II) is proved by Spitzer in [6, p. 327] (resp. p. 324).

EXAMPLE 2. Let \leq denote a lexicographic order on F and let J be the set H of all $f \in F^+$ such that $f = f' f''$ for $f', f'' \in F^+$ implies $f < f' f''$. Let $Y_h = \{ h \}$, for each $h \in H$. The fact that I, I' and II are satisfied is due to Chen, Fox and Lyndon [1] (cf. also [7]). A similar result holds when H is replaced by the set obtained by "removing the brackets" from Hall's *basic commutators* ([5] and [3, Chapter 11]).

We conclude with the following application of the "elimination method" of Lazard [4].

PROPOSITION 2. Let F be a free monoid, and P_1 and P_2 two subsets of F such that $F^+ = P_1 + P_2$. Then there exists a unique pair of subsets $Y_1 \subset P_1$ and $Y_2 \subset P_2$ such that

$$(4) \quad F = (1 - Y_1)^{-1}(1 - Y_2)^{-1}.$$

PROOF. Let X be a free set of generators of F and let $X_{i,0} = X \cap P_i$ ($i = 1, 2$). Then $W_0 = (1 - X_{2,0})^{-1}X_{2,0}X_{1,0}(1 - X_{1,0})^{-1}$ is the sum of all $f = f_2f_1$ where f_1 is a nontrivial word in the elements of $X_{1,0}$ and f_2 in those of $X_{2,0}$. It follows that $F = (1 - X_{1,0})^{-1}(1 - W_0)^{-1}(1 - X_{2,0})^{-1}$. If we let $Y_{i,0} = X_{i,0}$ ($i = 1, 2$) this establishes for $k=0$ the inductive hypothesis that

$$(5) \quad X_{i,k} \subset Y_{i,k} \subset P_i \quad (i = 1, 2) \quad \text{and} \quad F = F_{1,k}(1 - W_k)^{-1}F_{2,k}$$

where

$$F_{i,k} = (1 - Y_{i,k})^{-1} \quad (i = 1, 2) \quad \text{and} \quad W_k = F_{2,k}X_{2,k}X_{1,k}F_{1,k}.$$

Suppose (5) is satisfied for some $k \geq 0$. We construct inductively a sequence of subsets $W_{k,n}$ of W_k for all $n \geq 0$. First we take $W_{k,0} = \emptyset$. Supposing $W_{k,n}$ given we define $W_{k,n+1}$ to be the union of $W_{k,n}$ with the set of all words of minimal length in the complement of $(W_{k,n} \cap P_1)F_{1,k} \cup F_{2,k}(W_{k,n} \cap P_2)$ in W_k . We now define

$$X_{i,k+1} = \bigcup_{n \geq 0} (W_{k,n} \cap P_i); \quad Y_{i,k+1} = Y_{i,k} \cup X_{i,k+1} \quad (i = 1, 2).$$

Thus, $X_{i,k+1} \subset Y_{i,k+1} \subset P_i$ ($i = 1, 2$). To complete the verification that (5) holds for $k+1$, we need to show first

$$(6) \quad W_k = X_{1,k+1}F_{1,k} + F_{2,k}X_{2,k+1}.$$

Indeed, by the inductive hypothesis each $f \in W_k$ has a unique representation in the form $f = f_2f_1$, where $f_1 \in X_{1,k}F_{1,k}$ and $f_2 \in F_{2,k}X_{2,k}$. Taking $W_k = F_{2,k}W_kF_{1,k}$ into account, it follows that there exist two sets T_1 and T_2 such that $T_1 = X_{1,k+1}F_{1,k}$, $T_2 = F_{2,k}X_{2,k+1}$, and $W_k = T_1 \cup T_2$. Thus the proof of (6) needs only the verification that $T_1 \cap T_2 = \emptyset$.

Let $f \in T_2$. By definition $f = g_2f_2f_1$, where $g_2 \in F_{2,k}$, $f_2 \in F_{2,k}X_{2,k}$, $f_1 \in X_{1,k}F_{1,k}$, and $f_2f_1 \in W_{k,n} \cap P_1$ for some $n \geq 0$. The definition of $W_{k,n}$ implies that $f_2f_1 \notin X_{1,k+1}F_{1,k}$. Thus, for each $n' \geq 0$ and for each left factor $f'_1 \in X_{1,k}F_{1,k}$ of f_1 , we have $f_2f'_1 \notin W_{k,n'} \cap P_1$. It follows that for each such f'_1 we have $f_2f'_1 \in T_2$, hence $g_2f_2f'_1 \in T_2$, and finally $g_2f_2f'_1 \in W_{k,n''} \cap P_1$ for all $n'' \geq 0$. This shows that $f = g_2f_2f_1 \in T_1$ and $T_1 \cap T_2 = \emptyset$, hence (6) is proved.

For the rest, we compute as follows:

$$\begin{aligned}
& (F_{1,k+1}(1 - W_{k+1})^{-1}F_{2,k+1})^{-1} \\
&= (1 - Y_{2,k+1})(1 - (1 - Y_{2,k+1})^{-1}X_{2,k+1}X_{1,k+1}(1 - Y_{1,k+1})^{-1})(1 - Y_{1,k+1}) \\
&= 1 - Y_{2,k+1} - Y_{1,k+1} + Y_{2,k+1}Y_{1,k+1} - X_{2,k+1}X_{1,k+1} \\
&= 1 - (Y_{2,k} + X_{2,k+1}) - (Y_{1,k} + X_{1,k}) + (Y_{2,k} + X_{2,k+1})(Y_{1,k} + X_{1,k+1}) \\
&\quad - X_{2,k+1}X_{1,k+1} \\
&= 1 - Y_{2,k} - Y_{1,k} + Y_{2,k}Y_{1,k} - (1 - Y_{2,k})X_{1,k+1} - X_{2,k+1}(1 - Y_{1,k}) \\
&= (1 - Y_{2,k})(1 - X_{1,k+1}(1 - Y_{1,k})^{-1} - (1 - Y_{2,k})^{-1}X_{2,k+1})(1 - Y_{1,k}) \\
&= F_{2,k}^{-1}(1 - W_k)F_{1,k}^{-1} = F^{-1}
\end{aligned}$$

Finally, since $W_0 \subset FXXF$ and $W_{k+1} \subset FW_kW_kF$, each W_k ($k \geq 0$) contains no word of length less than 2^{k+1} . It follows that the same is true for the set complement of $(1 - Y_{1,k})^{-1}(1 - Y_{2,k})^{-1}$ in F . Thus, letting $Y_i = \bigcup_{k \geq 0} Y_{i,k}$ ($i = 1, 2$), we have proved the existence of at least one pair of sets satisfying the conditions stated in Proposition 2.

To verify the uniqueness, let us consider any other pair of subsets Y'_1 and Y'_2 of F^+ that satisfies $F = (1 - Y'_1)^{-1}(1 - Y'_2)^{-1}$. We define the subsets U_i, V_i, V'_i ($i = 1, 2$) of F^+ by the relations $U_i = Y_i \cap Y'_i$; $V_i = Y_i - U_i$; $V'_i = Y'_i - U_i$ ($i = 1, 2$). From $F^{-1} = (1 - Y_2)(1 - Y_1) = (1 - Y'_2)(1 - Y'_1)$ we deduce

$$(7) \quad -V_2 - V_1 + U_2V_1 + V_2U_1 = -V'_1 - V'_2 + U_2V'_1 + V'_2U_1.$$

Now, either $Y_1 = Y'_1$ and $Y_2 = Y'_2$ (i.e., $V_1 \cup V_2 \cup V'_1 \cup V'_2 = \emptyset$) or else $V_1 \cup V_2 \cup V'_1 \cup V'_2$ contains some element f of minimal length. By construction $V_1 \cap V'_1 = V_2 \cap V'_2 = \emptyset$. Thus (7) shows that $f \in (V_1 \cap V'_2) \cup (V_2 \cap V'_1)$. Since this last set is empty if $Y'_1 \subset P_1$ and $Y'_2 \subset P_2$, the verification of Proposition 2 is concluded.

Acknowledgment. I gratefully acknowledge the help of the referee in improving the proofs.

REFERENCES

1. K. T. Chen, R. H. Fox and R. C. Lyndon, *Free differential calculus*. IV, Ann. of Math. (2) **68** (1958), 81-95.
2. P. M. Cohn, *On subsemigroups of free semigroups*, Proc. Amer. Math. Soc. **13** (1962), 347-357.
3. Marshall Hall, *The theory of groups*, Macmillan, New York, 1959.
4. M. Lazard, *Groupes, anneaux de Lie et problème de Burnside*, Inst. Mat. dell'Università, Roma, 1960.
5. H. Meyer-Wunderli, *Note on a basis of P. Hall for the higher commutators in free groups*, Comment. Math. Helv. **26** (1952), 1-5.
6. F. Spitzer, *A combinatorial lemma and its application to probability theory*. Trans. Amer. Math. Soc., **82** (1956), 323-339.
7. A. I. Širšov, *On free Lie rings*, Mat. Sb. (N.S.) **45** (87) (1958), 113-122.

UNIVERSITÉ DE POITIERS, POITIERS, FRANCE