

COUNTING OCCURRENCES OF 132 IN AN EVEN PERMUTATION

Toufik Mansour ¹

Department of Mathematics, Chalmers University of Technology, S-41296 Göteborg,
Sweden

toufik@math.chalmers.se

ABSTRACT

We study the generating function for the number of even (or odd) permutations on n letters containing exactly $r \geq 0$ occurrences of a 132 pattern. It is shown that finding this function for a given r amounts to a routine check of all permutations in \mathfrak{S}_{2r} .

Nous tudions la fonction gnratrice pour le nombre de permutations paires (ou impaires) ayant n lettres contenant exactement $r \geq 0$ occurrences du motif 132. Nous montrons que trouver cette fonction pour r donn revient vrifier une routine pour toutes les permutations dans \mathfrak{S}_{2r} .

2000 Mathematics Subject Classification: Primary 05A05, 05A15; Secondary 05C90

1. INTRODUCTION

Let $[n] = \{1, 2, \dots, n\}$ and \mathfrak{S}_n denote the set of all permutations of $[n]$. We shall view permutations in \mathfrak{S}_n as words with n distinct letters in $[n]$. A pattern is a permutation $\sigma \in \mathfrak{S}_k$, and an occurrence of σ in a permutation $\pi = \pi_1\pi_2 \cdots \pi_n \in \mathfrak{S}_n$ is a subsequence of π that is order equivalent to σ . For example, an occurrence of 132 is a subsequence $\pi_i\pi_j\pi_k$ ($1 \leq i < j < k \leq n$) of π such that $\pi_i < \pi_k < \pi_j$. We denote by $\tau(\pi)$ the number of occurrences of τ in π , and we denote by $s_\sigma^r(n)$ the number of permutations $\pi \in \mathfrak{S}_n$ such that $\sigma(\pi) = r$.

In the last decade much attention has been paid to the problem of finding the numbers $s_\sigma^r(n)$ for a fixed $r \geq 0$ and a given pattern τ (see [1, 2, 3, 5, 6, 7, 10, 11, 14, 15, 16, 17, 18, 19]). Most of the authors consider only the case $r = 0$, thus studying permutations *avoiding* a given pattern. Only a few papers consider the case $r > 0$, usually restricting themselves to patterns of length 3. Using two simple involutions (*reverse* and *complement*) on \mathfrak{S}_n it is immediate that with respect to being equidistributed, the six patterns of length three fall into the two classes $\{123, 321\}$ and $\{132, 213, 231, 312\}$. Noonan [13] proved that

$$s_{123}^1(n) = \frac{3}{n} \binom{2n}{n-3}.$$

¹Research financed by EC's IHRP Programme, within the Research Training Network "Algebraic Combinatorics in Europe", grant HPRN-CT-2001-00272

A general approach to the problem was suggested by Noonan and Zeilberger [14]; they gave another proof of Noonan's result, and conjectured that

$$s_{123}^2(n) = \frac{59n^2 + 117n + 100}{2n(2n-1)(n+5)} \binom{2n}{n-4}$$

and

$$s_{132}^1(n) = \binom{2n-3}{n-3}.$$

The first conjecture was proved by Fulmek [8] and the latter conjecture was proved by Bóna in [6]. A conjecture of Noonan and Zeilberger states that $s_{\sigma}^r(n)$ is P -recursive in n for any r and τ . It was proved by Bóna [4] for $\sigma = 132$. Mansour and Vainshtein [11] suggested a new approach to this problem in the case $\sigma = 132$, which allows one to get an explicit expression for $s_{132}^r(n)$ for any given r . More precisely, they presented an algorithm that computes the generating function $\sum_{n \geq 0} s_{132}^r(n)x^n$ for any $r \geq 0$. To get the result for a given r , the algorithm performs certain routine checks for each element of the symmetric group \mathfrak{S}_{2r} . The algorithm has been implemented in C and yields explicit results for $1 \leq r \leq 6$.

Let π be any permutation. The number of *inversions* of π is given by $i_{\pi} = |\{(i, j) : \pi_i > \pi_j, i < j\}|$. The *signature* of π is given by $\text{sign}(\pi) = (-1)^{i_{\pi}}$. We say π is an *even permutation* (respectively; *odd permutation*) if $\text{sign}(\pi) = 1$ (respectively; $\text{sign}(\pi) = -1$). We denote by E_n (respectively; O_n) the set of all even (respectively; odd) permutations in \mathfrak{S}_n . Clearly, $|E_n| = |O_n| = \frac{1}{2}n!$ for all $n \geq 2$. The following lemma holds immediately by definitions.

Lemma 1.1. *For any permutation π , $\text{sign}(\pi) = (-1)^{21(\pi)}$.*

We denote by $e_{\sigma}^r(n)$ (respectively; $o_{\sigma}^r(n)$) the number of even (respectively; odd) permutations $\pi \in E_n$ (respectively; $\pi \in O_n$) such that $\sigma(\pi) = r$.

Apparently, for the first time the relation between even (odd) permutations and pattern avoidance problem was suggested by Simion and Schimidet in [16] for $\sigma \in \mathfrak{S}_3$. In particular, Simion and Schmidt [16] proved that

$$e_{132}^0(n) = \frac{1}{2(n+1)} \binom{2n}{n} + \frac{1}{n+1} \binom{n-1}{(n-1)/2},$$

and

$$o_{132}^0(n) = \frac{1}{2(n+1)} \binom{2n}{n} - \frac{1}{n+1} \binom{n-1}{(n-1)/2}.$$

In this paper, as a consequence of [12], we suggest a new approach to this problem in the case of even (or odd) permutations where $\sigma = 132$, which allows one to get an explicit expression for $e_{132}^r(n)$ for any given r . More precisely, we present an algorithm that computes the generating functions $E_r(x) = \sum_{n \geq 0} e_{132}^r(n)x^n$ and $O_r(x) = \sum_{n \geq 0} o_{132}^r(n)x^n$ for any $r \geq 0$. To get the result for a given r , the algorithm performs certain routine checks for each element of the symmetric group \mathfrak{S}_{2r} . The algorithm has been implemented in C, and yields explicit results for $0 \leq r \leq 6$.

2. DEFINITIONS AND PRELIMINARY RESULTS

In this section, we recall the definitions and the preliminary results of counting occurrences of the pattern 132 in a permutation, see [12]. Then, as a consequence of these results, we get the preliminary results of counting occurrences of the pattern 132 in an even (odd) permutation.

To any $\pi \in \mathfrak{S}_n$ we assign a bipartite graph G_π in the following way. The vertices in one part of G_π , denoted V_1 , are the entries of π , and the vertices of the second part, denoted V_3 , are the occurrences of 132 in π . Entry $i \in V_1$ is connected by an edge to occurrence $j \in V_3$ if i enters j . For example, let $\pi = 57614283$, then π contains 5 occurrences of 132, and the graph G_π is presented on Figure 1.




FIGURE 1. Graph G_π for $\pi = 57614283$

We denote by G_π^n the connected component of G_π containing the last entry n . Let $\pi(i_1), \dots, \pi(i_s)$ be the entries of π belonging to G_π^n , and let $\sigma = \sigma_\pi \in \mathfrak{S}_s$ be the corresponding pattern permutation. We say that $\pi(i_1), \dots, \pi(i_s)$ is the *kernel* of π and denote it $\ker \pi$ and σ is called the *shape* of the kernel, or the *kernel shape*. Let $s = s(\sigma_\pi)$ be the *size* of the kernel, and the number of occurrences of 132 in $\ker \pi$ is called the *capacity* of the kernel. For example, for $\pi = 57614283$ as above, the kernel equals 14283, its shape is 14253, the size equals 5, and the capacity equals 4.

Theorem 2.1. (Mansour and Vainshtein [12, Theorem 1]) *Let $\pi \in \mathfrak{S}_n$ such that $132(\pi) = r$, then the size of the kernel of π is at most $2r + 1$.*

We say that ρ is a *kernel permutation* if it is the kernel shape for some permutation π . Evidently ρ is a kernel permutation if and only if $\sigma_\rho = \rho$. Let $\rho \in \mathfrak{S}_s$ be an arbitrary kernel permutation. We denote by $\mathfrak{S}(\rho)$ the set of all the permutations of all possible sizes whose kernel shape equals ρ . For any $\pi \in \mathfrak{S}(\rho)$ we define the *kernel cell decomposition* as follows. The number of cells in the decomposition equals $s(s + 1)$. Let $\ker \pi = \pi(i_1), \dots, \pi(i_s)$; the cell $C_{ml} = C_{ml}(\pi)$ for $1 \leq l \leq s + 1$ and $1 \leq m \leq s$ is defined by $C_{ml}(\pi) = (\pi(j_1), \pi(j_2), \dots, \pi(j_d))$ such that $j_p \neq j_q$ if and only if $p \neq q$, and

$$\{j_1, j_2, \dots, j_d\} = \{j : i_{l-1} < j < i_l, \pi(i_{\rho^{-1}(m-1)}) < \pi(j) < \pi(i_{\rho^{-1}(m)})\},$$

where $i_0 = 0$, $i_{s+1} = n + 1$, $\rho^{-1}(0) = 0$, and $\pi(0) = 0$ for any π . If π coincides with ρ itself, then all the cells in the decomposition are empty. An arbitrary permutation in $\mathfrak{S}(\rho)$ is obtained by filling in some of the cells in the cell decomposition. A cell C is called *infeasible* if the existence of an entry $a \in C$ would imply an occurrence of 132 that contains a and two other entries $x, y \in \ker \pi'$ for some π' . Clearly, all infeasible cells are empty for any $\pi \in \mathfrak{S}(\rho)$. All the remaining cells are called *feasible*; a feasible cell may, or may not, be empty. Consider the permutation $\pi = 67382451$. The kernel

of π equals 3845, its shape is 1423. The cell decomposition of π contains four feasible cells: $C_{13} = (2)$, $C_{14} = \emptyset$, $C_{15} = (1)$, and $C_{41} = (6, 7)$ (see Figure 2). All the other cells are infeasible; for example, C_{32} is infeasible, since if $a \in C_{32}$, then $a\pi'(i_2)\pi'(i_4)$ is an occurrence of 132 for some $\ker \pi'$.




FIGURE 2. Kernel cell decomposition for $\pi = 67382451 \in \mathfrak{S}(1423)$

As another example, permutation $\tilde{\pi} = 11\ 10\ 7\ 12\ 4\ 6\ 5\ 8\ 2\ 9\ 3\ 1$ belongs to $\mathfrak{S}(364152)$. Its kernel is $7\ 12\ 8\ 2\ 9\ 3$, and the feasible cells are $C_{33} = (4, 6, 5)$, $C_{17} = (1)$, $C_{61} = (11, 10)$.

Given a cell C_{ij} in the kernel cell decomposition, all the kernel entries can be positioned with respect to C_{ij} . We say that $x = \pi(i_k) \in \ker \pi$ lies *below* C_{ij} if $\rho(k) < i$, and *above* C_{ij} if $\rho(k) \geq i$. Similarly, x lies to the *left* of C_{ij} if $k < j$, and to the *right* of C_{ij} if $k \geq j$. As usual, we say that x lies to the *southwest* of C_{ij} if it lies below C_{ij} and to the left of it; the other three directions, northwest, southeast, and northeast, are defined similarly. Let us define a partial order \prec on the set of all feasible cells by saying that $C_{ml} \prec C_{m'l'} \neq C_{ml}$ if $m \geq m'$ and $l \leq l'$.

Theorem 2.2. (Mansour and Vainshtein [12])

- (i) \prec is a linear order.
- (ii) Let C_{ml} and $C_{m'l'}$ be two nonempty feasibly cells such that $l < l'$. Then for any pair of entries $a \in C_{ml}$, $b \in C_{m'l'}$, one has $a > b$.
- (iii) Let C_{ml} and $C_{m'l}$ be two nonempty feasibly cells such that $m < m'$. Then any entry $a \in C_{ml}$ lies to the right of any entry $b \in C_{m'l}$ ($\pi^{-1}(b) < \pi^{-1}(a)$).

Let π be any permutation with a kernel permutation ρ , and assume that the feasible cells of the kernel cell decomposition associated with ρ are ordered linearly according to \prec , $C^1, C^2, \dots, C^{f(\rho)}$. Let d_j be the size of C^j . For example, let $\pi = 67382451$ with kernel permutation $\rho = 1423$, as on Figure 2, then $d_1 = 2$, $d_2 = 1$, $d_3 = 0$, and $d_4 = 1$. We denote by $l_j(\rho)$ the number of the entries of ρ that lie to the north-west from C^j or lie to the south-east from C^j . For example, let $\rho = 1423$, as on Figure 2, then $l_1(\rho) = 3$, $l_2(\rho) = 2$, $l_3(\rho) = 3$, and $l_4(\rho) = 4$. Clearly, $l_1(\rho) = s(\rho) - 1$ and $l_{f(\rho)} = s(\rho)$ for any nonempty kernel permutation ρ . We denote by $\text{sign}(C^j)$ the number of occurrences of a 21 pattern in C^j .

Lemma 2.3. *For any permutation π with a kernel permutation ρ ,*

$$\text{sign}(\pi) = (-1)^{\left(\sum_{1 \leq i \leq j \leq f(\rho)} d_i d_j + \sum_{j=1}^{f(\rho)} d_j l_j(\rho)\right)} \cdot \text{sign}(\rho) \cdot \prod_{j=1}^{f(\rho)} \text{sign}(C^j).$$

Proof. To verify this formula, let us count the number of occurrences of the pattern 21 in π . There four possibilities for an occurrence of 21 in π . The first possibility is an occurrence occurs in one of the cells C^j , so in this case there are $\sum_{j=1}^{f(\rho)} 21(C^j)$ occurrences. The second possibility is an occurrence occurs in $\ker \pi$, so there are $21(\rho)$ occurrences. The third possibility is an occurrence of two elements which the first belongs to $\ker \pi$ and the second belongs to C^i , so there are $\sum_{j=1}^{f(\rho)} d_j l_j(\rho)$ (see Theorem 2.2) occurrences. The fourth possibility is an occurrence of two elements which the first belongs to C^i and the second belongs to C^j where $i < j$ (Theorem 2.2 yields every entry of C^i is greater than every entry of C^j for all $i < j$), so there are $\sum_{1 \leq i < j \leq f(\rho)} d_i d_j$ occurrences. Therefore, by Lemma 1.1 we have

$$\text{sign}(\pi) = (-1)^{\sum_{j=1}^{f(\rho)} 21(C^j)} (-1)^{21(\rho)} (-1)^{\sum_{j=1}^{f(\rho)} d_j l_j(\rho)} (-1)^{\sum_{1 \leq i < j \leq f(\rho)} d_i d_j},$$

equivalently, $\text{sign}(\pi) = (-1)^{\left(\sum_{1 \leq i \leq j \leq f(\rho)} d_i d_j + \sum_{j=1}^{f(\rho)} d_j l_j(\rho)\right)} \cdot \text{sign}(\rho) \cdot \prod_{j=1}^{f(\rho)} \text{sign}(C^j)$. \square

We say the vector $\mathbf{v} = (v_1, v_2, \dots, v_n)$ is a *binary vector* if $v_i \in \{0, 1\}$ for all i , $1 \leq i \leq n$. We denote the set of all binary vectors of length n by \mathcal{B}^n . For any $\mathbf{v} \in \mathcal{B}^n$, we define $|\mathbf{v}| = v_1 + v_2 + \dots + v_n$. For example, $\mathcal{B}^2 = \{(0, 0), (0, 1), (1, 0), (1, 1)\}$ and $|(1, 1, 0, 0, 1)| = 3$.

Let ρ be any kernel permutations; we denote by $X_a(\rho)$ (respectively; Y_a^ρ) the set of all the binary vectors $\mathbf{v} \in \mathcal{B}^{f(\rho)}$ such that $(-1)^{|\mathbf{v}|+s(\rho)} = a$ (respectively; $(-1)^{|\mathbf{v}|} = a$). For any $\mathbf{v} \in \mathcal{B}^{f(\rho)}$, we define

$$z_\rho(\mathbf{v}) = (-1)^{\sum_{1 \leq i < j \leq f(\rho)} v_i v_j + \sum_{j=1}^{f(\rho)} l_j(\rho) v_j} \text{sign}(\rho).$$

Letting ρ be any kernel permutations and $\mathbf{v} = (v_1, v_2, \dots, v_{f(\rho)}) \in \mathcal{B}^{f(\rho)}$, we denote by $\mathfrak{S}(\rho; \mathbf{v})$ the set of all permutations of all sizes with kernel permutation ρ such that the corresponding cells C^j satisfy $(-1)^{d_j} = (-1)^{v_j}$; in such a context \mathbf{v} is called a *length argument vector* of ρ . By definitions, the following result holds immediately.

Lemma 2.4. *For any kernel permutation ρ ,*

$$\mathfrak{S}(\rho) = \bigcup_{\mathbf{v} \in \mathcal{B}^{f(\rho)}} \mathfrak{S}(\rho; \mathbf{v}).$$

Letting ρ be any kernel permutations and letting

$$\mathbf{v} = (v_1, v_2, \dots, v_{f(\rho)}), \mathbf{u} = (u_1, u_2, \dots, u_{f(\rho)}) \in \mathcal{B}^{f(\rho)},$$

we denote by $\mathfrak{S}(\rho; \mathbf{v}, \mathbf{u})$ the set of all permutations in $\mathfrak{S}(\rho; \mathbf{v})$ such that the corresponding cells C^j satisfy $\text{sign}(C^j) = 1$ if and only if $u_j = 0$; in such a context \mathbf{u} is called a *signature argument vector* of ρ . By Lemma 2.4, the following result holds immediately.

Lemma 2.5. *For any kernel permutation ρ ,*

$$\mathfrak{S}(\rho) = \bigcup_{\mathbf{v} \in \mathcal{B}^{f(\rho)}} \mathfrak{S}(\rho; \mathbf{v}) = \bigcup_{\mathbf{v} \in \mathcal{B}^{f(\rho)}} \bigcup_{\mathbf{u} \in \mathcal{B}^{f(\rho)}} \mathfrak{S}(\rho; \mathbf{v}, \mathbf{u}).$$

For any $a, b \in \{0, 1\}$ we define

$$H_r(a, b) = \begin{cases} \frac{1}{2}(E_r(x) + (-1)^a E_r(-x)), & \text{if } b = 0 \\ \frac{1}{2}(O_r(x) + (-1)^a O_r(-x)), & \text{if } b = 1 \end{cases}.$$

By definitions, the following result holds immediately.

Lemma 2.6. *Let $a, b \in \{0, 1\}$. Then the generating function for all permutations π such that $132(\pi) = r$, $(-1)^{|\pi|} = (-1)^a$, and $\text{sign}(\pi) = (-1)^b$ is given by $H_r(a, b)$, where we denote by $|\pi|$ the length of the permutation π .*

3. MAIN THEOREM

The main result of this note can be formulated as follows. Denote by K the set of all kernel permutations, and by K_t the set of all kernel shapes for permutations in \mathfrak{S}_t . Letting ρ be any kernel permutation, for any $a, b \in \{0, 1\}$ and any $r_1, \dots, r_{f(\rho)}$, we define

$$L_{r_1, \dots, r_{f(\rho)}}^\rho(a, b) = \sum_{\mathbf{v} \in X_{(-1)^a}^\rho} \sum_{\mathbf{u} \in Y_{(-1)^b z_\rho(\mathbf{v})}^\rho} \prod_{j=1}^{f(\rho)} H_{r_j}(v_j, u_j).$$

Remark 3.1. *The product $\prod_{j=1}^s H_{r_j}(v_j, u_j)$ gives the generating function for the number of s permutations π^1, \dots, π^s such that π^j contains 132 exactly r_j times, the modulo 2 of the length of π^j is v_j , and π^j is even (respectively; odd) permutation if and only if $u_j = 1$ (respectively; $u_j = 0$).*

Theorem 3.2. *Let $r \geq 1$. For any $a, b \in \{0, 1\}$,*

$$(3.1) \quad H_r(a, b) = \sum_{\rho \in K_{2r+1}} \sum_{r_1 + \dots + r_{f(\rho)} = r - c(\rho), r_j \geq 0} L_{r_1, \dots, r_{f(\rho)}}^\rho(a, b).$$

Proof. Let us fix a kernel permutation $\rho \in K_{2r+1}$, a length argument vector $\mathbf{v} = (v_1, \dots, v_{f(\rho)}) \in X_{(-1)^a}(\rho)$, and a signature argument vector $\mathbf{u} = (u_1, \dots, u_{f(\rho)}) \in Y_{(-1)^b z_\rho(\mathbf{v})}^{f(\rho)}$. Recall that the kernel ρ of any π contains exactly $c(\rho)$ occurrences of 132. The remaining $r - c(\rho)$ occurrences of 132 are distributed among the feasible cells of the kernel cell decomposition of π . By Theorem 2.2, each occurrence of 132 belongs entirely to one feasible cell, and the occurrences of 132 in different cells do not influence one another.

Let π be any permutation such that $132(\pi) = r$, $\text{sign}(\pi) = (-1)^b$ and $(-1)^{|\pi|} = (-1)^a$ together with a kernel permutations ρ , length argument vector \mathbf{v} , and signature argument vector \mathbf{u} . Then by Lemma 2.5, the cells C^j satisfy the following conditions:

- (1) $v_j = 0$ if and only if d_j is an even number,
- (2) $u_j = 0$ if and only if $\text{sign}(C^j) = 1$,
- (3) $(-1)^{v_1 + \dots + v_{f(\rho)} + s(\rho)} = (-1)^a$, and
- (4) $(-1)^{u_1 + \dots + u_{f(\rho)}} z_\rho(\mathbf{v}) = (-1)^b$.

Therefore, by Remark ?? and Lemma 2.6 this contribution gives

$$x^{s(\rho)} \sum_{r_1 + \dots + r_{f(\rho)} = r - c(\rho), r_j \geq 0} \prod_{j=1}^{f(\rho)} H_{r_j}(v_j, u_j).$$

Hence by Lemma 2.5 and Theorem 2.1, if summing over all the kernel permutations $\rho \in K_{2r+1}$, length argument vectors $\mathbf{v} \in X_{(-1)^a}(\rho)$, and signature argument vectors $\mathbf{u} \in Y_{(-1)^b}^{f(\rho)} z_\rho(\mathbf{v})$ then we get the desired result. \square

Proposition 3.3. (see Mansour and Vainshtein [12, Proposition]) *The only kernel permutation of capacity $r \geq 1$ and size $2r + 1$ is*

$$\rho = 2r - 1, 2r + 1, 2r - 3, 2r, \dots, 2r - 2j - 3, 2r - 2j, \dots, 1, 4, 2.$$

Theorem 3.2 provides a finite algorithm for finding $E_r(x)$ and $O_r(x)$ for any given $r \geq 0$, since we only have to consider all permutations in \mathfrak{S}_{2r+1} , and to perform certain routine operations with all shapes found so far. Moreover, the amount of search can be decreased substantially due to the following theorem.

Theorem 3.4. *The only kernel permutation of capacity $r \geq 1$ and size $2r + 1$ is*

$$\rho = 2r - 1, 2r + 1, 2r - 3, 2r, \dots, 2r - 2j - 3, 2r - 2j, \dots, 1, 4, 2.$$

Its parameters are given by $s(\rho) = 2r + 1$, $c(\rho) = r$, $f(\rho) = r + 2$, $\text{sign}(\rho) = -1$, and $z_\rho(v_1, \dots, v_{r+2}) = (-1)^{(1+v_{r+2} + \sum_{1 \leq i < j \leq r+2} v_i v_j)}$.

Proof. The first part of the proposition holds by Proposition 3.3. Besides, by using the form of ρ we get $s(\rho) = 2r + 1$, $c(\rho) = r$, $f(\rho) = r + 2$; $\text{sign}(\rho) = -1$, and $l_j(\rho) = 2r$ for all $j = 1, 2, \dots, r + 1$ and $l_{r+2}(\rho) = 2r + 1$. Therefore, $z_\rho(v_1, \dots, v_{r+2}) = (-1)^{(1+v_{r+2} + \sum_{1 \leq i < j \leq r+2} v_i v_j)}$. \square

By this proposition, it suffices to search only permutations in \mathfrak{S}_{2r} . Below we present several explicit calculations.

3.1. The case $r = 0$. Let us start from the case $r = 0$. Observe that Theorem 3.2 remains valid for $r = 0$, provided the left hand side of Equation 3.1 for $a = b = 0$ is replaced by $H_r(0, 0) - 1 = \frac{1}{2}(E_r(x) + E_r(-x)) - 1$; subtracting 1 here accounts for the empty permutation. So, we begin with finding kernel shapes for all permutations in \mathfrak{S}_1 . The only shape obtained is $\rho_1 = 1$, and it is easy to see that $s(\rho_1) = 1$, $c(\rho_1) = 0$, $f(\rho_1) = 2$,

$$X_1(\rho_1) = Y_{-1} = \{(1, 0), (0, 1)\}, \quad X_{-1}(\rho_1) = Y_1 = \{(0, 0), (1, 1)\},$$

and

$$z_{\rho_1}(0, 0) = z_{\rho_1}(1, 0) = z_{\rho_1}(1, 1) = -z_{\rho_1}(0, 1) = 1.$$

Therefore, Equation 3.1 for $a = b = 0$ gives

$$(3.2) \quad \begin{aligned} & \frac{1}{2}(E_0(x) + E_0(-x)) - 1 = \\ & = xH_0(1, 0)H_0(0, 0) + xH_0(1, 1)H_0(0, 1) + xH_0(1, 0)H_0(0, 1) + xH_0(1, 1)H_0(0, 0), \end{aligned}$$

Equation 3.1 for $a = 1$ and $b = 0$ gives

$$(3.3) \quad \frac{1}{2}(E_0(x) - E_0(-x)) = xH_0^2(0, 0) + xH_0^2(0, 1) + xH_0^2(1, 0) + xH_0^2(1, 1),$$

Equation 3.1 for $a = 0$ and $b = 1$ gives

$$(3.4) \quad \begin{aligned} & \frac{1}{2}(O_0(x) + O_0(-x)) = \\ & = xH_0(1, 1)H_0(0, 0) + xH_0(1, 0)H_0(0, 1) + xH_0(0, 0)H_0(1, 0) + xH_0(0, 1)H_0(1, 1), \end{aligned}$$

and Equation 3.1 for $a = b = 1$ gives

$$(3.5) \quad \frac{1}{2}(O_0(x) - O_0(-x)) = 2xH_0(0, 1)H_0(0, 0) + 2xH_0(1, 1)H_0(1, 0).$$

Our present aim to find explicitly $E_0(x)$ and $O_0(x)$, thus we need the following notation. We define

$$M_r(x) = E_r(x) - O_r(x) \text{ and } F_r(x) = E_r(x) + O_r(x)$$

for all $r \geq 0$. Clearly,

$$\begin{aligned} H_r(0, 0) - H_r(0, 1) &= \frac{1}{2}(M_r(x) + M_r(-x)), & H_r(0, 0) + H_r(0, 1) &= \frac{1}{2}(F_r(x) + F_r(-x)), \\ H_r(1, 0) - H_r(1, 1) &= \frac{1}{2}(M_r(x) - M_r(-x)), & H_r(1, 0) + H_r(1, 1) &= \frac{1}{2}(F_r(x) - F_r(-x)), \end{aligned}$$

for all $r \geq 0$. Therefore, by subtracting (respectively; adding) Equation 3.4 and Equation 3.2, and by subtracting (respectively; adding) Equation 3.5 and Equation 3.3 we get

$$\begin{cases} M_0(x) + M_0(-x) = 2 \\ M_0(x) - M_0(-x) = x(M_0^2(x) + M_0^2(-x)) \end{cases}$$

and

$$\begin{cases} F_0(x) + F_0(-x) = 2 + x(F_0^2(x) - F_0^2(-x)) \\ F_0(x) - F_0(-x) = x(F_0^2(x) + F_0^2(-x)). \end{cases}$$

Hence,

$$M_0(x) = 1 + \frac{1 - \sqrt{1 - 4x^2}}{2x} \text{ and } F_0(x) = \frac{1 - \sqrt{1 - 4x}}{2x}.$$

Theorem 3.5. (i) *The generating function for the number of even permutations avoiding 132 is given by (see [16])*

$$E_0(x) = \frac{1}{2} \left(\frac{1 - \sqrt{1 - 4x}}{2x} + 1 + \frac{1 - \sqrt{1 - 4x^2}}{2x} \right).$$

(ii) *The generating function for the number of odd permutations avoiding 132 is given by (see [16])*

$$O_0(x) = \frac{1}{2} \left(\frac{1 - \sqrt{1 - 4x}}{2x} - 1 - \frac{1 - \sqrt{1 - 4x^2}}{2x} \right).$$

(iii) *The generating function for the number of permutations avoiding 132 is given by (see [9])*

$$F_0(x) = \frac{1 - \sqrt{1 - 4x}}{2x}.$$

3.2. The Case $r = 1$. Since permutations in \mathfrak{S}_2 do not exhibit kernel shapes distinct from ρ_1 , the only possible new shape is the exceptional one, $\rho_2 = 132$. Calculation of the parameters of ρ_2 gives $s(\rho_2) = 3$, $c(\rho_2) = 1$, $f(\rho_2) = 3$,

$$\begin{aligned} X_1(\rho_2) &= Y_{-1} = \{(1, 0, 0), (0, 1, 0), (0, 0, 1), (1, 1, 1)\}, \\ X_{-1}(\rho_2) &= Y_1 = \{(0, 0, 0), (1, 1, 0), (1, 0, 1), (1, 1, 0)\}, \end{aligned}$$

and

$$\begin{aligned} z_{\rho_2}(0, 0, 0) &= z_{\rho_2}(1, 0, 0) = z_{\rho_2}(0, 1, 0) = -z_{\rho_2}(1, 1, 0) = 1, \\ -z_{\rho_2}(0, 0, 1) &= z_{\rho_2}(1, 0, 1) = z_{\rho_2}(0, 1, 1) = z_{\rho_2}(1, 1, 1) = 1. \end{aligned}$$

Therefore, by Theorem 3.2 we have

$$\left\{ \begin{array}{l} 2(H_1(0, 0) - H_1(0, 1)) = \\ \quad = M_1(x) + M_1(-x) = \frac{x^3}{2}(M_0(-x) - M_0(x))(M_0^2(-x) + M_0^2(x)) \\ \\ 2(H_1(1, 0) - H_1(1, 1)) = \\ \quad = M_1(x) - M_1(-x) = 2x(M_0(x)M_1(x) + M_0(-x)M_1(-x)) \\ \quad \quad - \frac{x^3}{2}(M_0(-x) + M_0(x))(M_0^2(-x) + M_0^2(x)), \end{array} \right.$$

Using the expression for $M_0(x)$ (see the case $r = 0$) we get

$$M_1(x) = \frac{1}{2}(-1 + 3x + 2x^2) + \frac{1 - 3x - 4x^2 + 4x^3}{2}(1 - 4x^2)^{-1/2}.$$

Similarly, if considering the expressions for $H_1(0, 0) + H_1(0, 1)$ and $H_1(1, 0) + H_1(1, 1)$ we get

$$F_1(x) = \frac{1}{2}(x - 1) + \frac{1 - 3x}{2}(1 - 4x)^{-1/2}.$$

Theorem 3.6. (i) *The generating function for the number of even permutations containing 132 exactly once is given by*

$$E_1(x) = -\frac{1}{2}(1 - 2x - x^2) + \frac{1 - 3x}{4}(1 - 4x)^{-1/2} + \frac{1 - 3x - 4x^2 + 4x^3}{4}(1 - 4x^2)^{-1/2}.$$

(ii) *The generating function for the number of odd permutations containing 132 exactly once is given by*

$$O_1(x) = -\frac{1}{2}(x + x^2) + \frac{1 - 3x}{4}(1 - 4x)^{-1/2} - \frac{1 - 3x - 4x^2 + 4x^3}{4}(1 - 4x^2)^{-1/2}.$$

(iii) *The generating function for the number of permutations containing 132 exactly once is given by (see [6])*

$$F_1(x) = \frac{1}{2}(x - 1) + \frac{1 - 3x}{2}(1 - 4x)^{-1/2}.$$

3.3. The case $r = 2$. We have to check the kernel shapes of permutations in \mathfrak{S}_4 . Exhaustive search adds four new shapes to the previous list; these are 1243, 1342, 1423, and 2143; besides, there is the exceptional $35142 \in \mathfrak{S}_5$. Calculation of the parameters s, c, f, z, X_a, Y_a is straightforward, and we get

Theorem 3.7. (i) *The generating function for the number of even permutations containing 132 exactly twice is given by*

$$E_2(x) = \frac{1}{2}x(x^3 + 3x^2 - 4x - 1) + \frac{1}{4}(2x^4 - 4x^3 + 29x^2 - 15x + 2)(1 - 4x)^{-3/2} \\ - \frac{1}{4}(16x^7 - 48x^6 - 76x^5 + 64x^4 + 36x^3 - 21x^2 - 5x + 2)(1 - 4x^2)^{-3/2}.$$

(ii) *The generating function for the number of odd permutations containing 132 exactly once is given by*

$$O_2(x) = -\frac{1}{2}(x^4 + 3x^3 - 5x^2 - 4x + 2) + \frac{1}{4}(2x^4 - 4x^3 + 29x^2 - 15x + 2)(1 - 4x)^{-3/2} \\ + \frac{1}{4}(16x^7 - 48x^6 - 76x^5 + 64x^4 + 36x^3 - 21x^2 - 5x + 2)(1 - 4x^2)^{-3/2}.$$

(iii) *The generating function for the number of permutations containing 132 exactly twice is given by (see [12])*

$$F_2(x) = \frac{1}{2}(x^2 + 3x - 2) + \frac{1}{2}(2x^4 - 4x^3 + 29x^2 - 15x + 2)(1 - 4x)^{-3/2}.$$

3.4. The cases $r = 3, 4, 5, 6$. Let $r = 3, 4, 5, 6$; exhaustive search in $\mathfrak{S}_6, \mathfrak{S}_8, \mathfrak{S}_{10}$, and \mathfrak{S}_{12} reveals 20, 104, 503, and 2576 new nonexceptional kernel shapes, respectively, and we get

Theorem 3.8. *Let $r = 3, 4, 5, 6$, then*

$$M_r(x) = \frac{1}{2} (A_r(x) + B_r(x)(1 - 4x^2)^{-r+1/2})$$

and

$$F_r(x) = \frac{1}{2} (C_r(x) + D_r(x)(1 - 4x)^{-r+1/2}),$$

where

$$A_3(x) = 2x^6 + 10x^5 - 24x^4 - 30x^3 + 23x^2 + 7x - 2,$$

$$A_4(x) = 2x^8 + 14x^7 - 46x^6 - 90x^5 + 117x^4 + 85x^3 - 42x^2 - 8x + 1,$$

$$A_5(x) = 2x^{10} + 18x^9 - 76x^8 - 198x^7 + 360x^6 + 440x^5 - 355x^4 - 171x^3 + 62x^2 \\ + 10x - 2,$$

$$A_6(x) = 256x^{13} - 446x^{12} - 618x^{11} + 194x^{10} - 140x^9 + 798x^8 + 1404x^7 - 1702x^6 \\ - 1430x^5 + 815x^4 + 302x^3 - 88x^2 - 15x + 4,$$

$$B_3(x) = 64x^{11} - 320x^{10} - 800x^9 + 1216x^8 + 1124x^7 - 972x^6 - 524x^5 + 312x^4 \\ + 100x^3 - 43x^2 - 7x + 2,$$

$$B_4(x) = -256x^{15} + 1792x^{14} + 6112x^{13} - 13120x^{12} - 19840x^{11} + 22224x^{10} \\ + 19054x^9 - 14780x^8 - 8328x^7 + 4772x^6 + 1840x^5 - 775x^4 - 197x^3 + 56x^2 \\ + 8x - 1,$$

$$B_5(x) = 1024x^{19} - 9216x^{18} - 40064x^{17} + 111744x^{16} + 228896x^{15} - 343264x^{14} \\ - 404056x^{13} + 398712x^{12} + 321058x^{11} - 234686x^{10} - 137468x^9 + 78480x^8 \\ + 33896x^7 - 15400x^6 - 4780x^5 + 1723x^4 + 351x^3 - 98x^2 - 10x + 2,$$

$$B_6(x) = 524288x^{24} + 1175552x^{23} - 1593344x^{22} - 2324992x^{21} + 1162752x^{20} \\ + 298112x^{19} + 2696448x^{18} + 4856864x^{17} - 7020288x^{16} - 7464568x^{15} \\ + 6981056x^{14} + 5445696x^{13} - 3868942x^{12} - 2335450x^{11} + 1324884x^{10} \\ + 627306x^9 - 290536x^8 - 106510x^7 + 40772x^6 + 11046x^5 - 3543x^4 \\ - 632x^3 + 176x^2 + 15x - 4,$$

$$C_3(x) = 2x^3 - 5x^2 + 7x - 2,$$

$$C_4(x) = 5x^4 - 7x^3 + 2x^2 + 8x - 3,$$

$$C_5(x) = 14x^5 - 17x^4 + x^3 - 16x^2 + 14x - 2,$$

$$C_6(x) = 42x^6 - 44x^5 + 5x^4 + 4x^3 - 20x^2 + 19x - 4,$$

and

$$D_3(x) = -22x^6 - 106x^5 + 292x^4 - 302x^3 + 135x^2 - 27x + 2,$$

$$D_4(x) = 2x^9 + 218x^8 + 1074x^7 - 1754x^6 + 388x^5 + 1087x^4,$$

$$D_5(x) = -50x^{11} - 2568x^{10} - 10826x^9 + 16252x^8 - 12466x^7 + 16184x^6 - 16480x^5 \\ + 9191x^4 - 2893x^3 + 520x^2 - 50x + 2,$$

$$D_6(x) = 4x^{14} + 820x^{13} + 32824x^{12} + 112328x^{11} - 205530x^{10} + 141294x^9 \\ - 30562x^8 - 67602x^7 + 104256x^6 - 74090x^5 + 30839x^4 - 7902x^3 + 1230x^2 \\ - 107x + 4.$$

Moreover, for $r = 3, 4, 5, 6$,

$$E_r(x) = \frac{1}{4} (A_r(x) + C_r(x) + D_r(x)(1 - 4x)^{-r+1/2} + B_r(x)(1 - 4x^2)^{-r+1/2})$$

and

$$O_r(x) = \frac{1}{4} (A_r(x) - C_r(x) + D_r(x)(1 - 4x)^{-r+1/2} - B_r(x)(1 - 4x^2)^{-r+1/2}).$$

4. FURTHER RESULTS AND OPEN QUESTIONS

First of all, let us simplify the expression

$$L_{r_1, \dots, r_{f(\rho)}}^{\rho}(a, 0) - L_{r_1, \dots, r_{f(\rho)}}^{\rho}(a, 1),$$

where $a = 0, 1$, $r_j \geq 0$ for all j .

Lemma 4.1. *Let $\mathbf{v} \in \{0, 1\}^n$ be any vector, and let $a \in \{1, -1\}$. Then*

$$\sum_{\mathbf{x} \in Y_a} \prod_{j=1}^n H_r(v_j, x_j) - \sum_{\mathbf{y} \in Y_{-a}} \prod_{j=1}^n H_r(v_j, y_j) = a \prod_{j=1}^n g_r(j),$$

where $g_r(j) = H_r(v_j, 0) - H_r(v_j, 1) = \frac{1}{2}(M_r(x) + (-1)^{v_j} M_r(-x))$ for all j .

Proof. Let us define an order on the set \mathcal{B}^n , we say the vector $\mathbf{v} < \mathbf{u}$ if there exists j such that $u_j + v_j = 1$, and $u_i = v_i$ for all $i \neq j$. We say the 2^n vectors $\mathbf{u}^1, \dots, \mathbf{u}^{2^n}$ of \mathcal{B}^n are satisfy the ℓ -property if

$$\mathbf{0} = (0, 0, \dots, 0) = \mathbf{u}^1 < \mathbf{u}^2 < \dots < \mathbf{u}^{2^n},$$

and we say the 2^n vectors $\mathbf{u}^1, \dots, \mathbf{u}^{2^n}$ are satisfy the c -property if the vectors

$$(\mathbf{u}_1^1, \dots, \mathbf{u}_m^1), \dots, (\mathbf{u}_1^{2^m}, \dots, \mathbf{u}_m^{2^m})$$

are satisfy the ℓ -property for all $m = 1, 2, \dots, n$. For example, the vectors of \mathcal{B}^3 are satisfy the c -property by

$$(0, 0, 0) < (1, 0, 0) < (1, 1, 0) < (0, 1, 0) < (0, 1, 1) < (1, 1, 1) < (1, 0, 1) < (0, 0, 1).$$

First of all, let us prove by induction on n that there exists an order of the vectors of \mathcal{B}^n with c -property, for all $n \geq 1$. For $n = 1$ the c -property holds with $(0) < (1)$. Suppose that there exists an order of the vector of \mathcal{B}^m with the c -property. Let $\mathbf{v}^j = (\mathbf{u}_1^j, \dots, \mathbf{u}_m^j, 0)$ for all $j = 1, 2, \dots, 2^m$ and $\mathbf{v}^{2^m+j} = (\mathbf{u}_1^{2^m+1-j}, \dots, \mathbf{u}_m^{2^m+1-j}, 1)$ for all $j = 1, 2, \dots, 2^m$. By definitions, $\mathbf{v}^1 = (0, 0, \dots, 0)$ and $\mathbf{v}^1 < \dots < \mathbf{v}^{2^m+1}$, so the ℓ -property holds for $m+1$. Hence, by induction on m we get that there exists an order of the vector of \mathcal{B}^n with the c -property.

Now we are ready to prove the lemma. Without loss of generality we can assume that $(0, 0, \dots, 0) \in Y_a^n$ (which means $a = 1$); otherwise it is enough to replace a by $-a$. Let $\mathbf{x}^1, \dots, \mathbf{x}^{2^n}$ all the vectors of \mathcal{B}^n with the c -property. Using $(0, 0, \dots, 0) \in Y_a^n$ together with the c -property we get that $\mathbf{x}^{2^{i-1}} \in Y_a^n$ and $\mathbf{x}^{2^i} \in Y_{-a}^n$ for all $i = 1, 2, \dots, 2^{n-1}$. Therefore, for all $i = 1, 2, \dots, 2^{n-1}$,

$$\begin{aligned} & \sum_{i=1}^{2^{n-1}} \left(\prod_{j=1}^n H_r(v_j, \mathbf{x}_j^{2^{i-1}}) - \prod_{j=1}^n H_r(v_j, \mathbf{x}_j^{2^i}) \right) \\ &= \sum_{i=1}^{2^{n-1}} \left((-1)^{i-1} g_r(1) \prod_{j=2}^n H_r(v_j, \mathbf{x}_j^{2^{i-1}}) \right) \\ &= g_r(1) \sum_{i=1}^{2^{n-2}} \left(\prod_{j=1}^{n-1} H_r(\tilde{v}_j, \mathbf{y}_j^{2^{i-1}}) - \prod_{j=1}^{n-1} H_r(\tilde{v}_j, \mathbf{y}_j^{2^i}) \right), \end{aligned}$$

where $\mathbf{y}^p = (\mathbf{x}_2^{2p}, \dots, \mathbf{x}_n^{2p})$ for all $p = 1, 2, \dots, 2^{n-1}$, and $\tilde{v} = (v_2, v_3, \dots, v_n)$. Using the c-property for $\mathbf{x}^1, \dots, \mathbf{x}^{2^n}$ we get that the vectors $\mathbf{y}^1, \dots, \mathbf{y}^{2^{n-1}}$ are satisfy the c-property in \mathcal{B}^{n-1} . Hence by induction on n (by definitions, the lemma holds for $n = 1$), we get that the expression equals to $a \prod_{j=1}^n g_r(j)$. \square

As a remark, the vector $(0, 0, \dots, 0) \in Y_{z_\rho(\mathbf{v})}^\rho$ if and only if $z_\rho(\mathbf{v}) = 1$ for any kernel permutation ρ and vector \mathbf{v} . Therefore, by Theorem 3.2 and Lemma 4.1 we get the following result.

Theorem 4.2. *Let $a \in \{0, 1\}$ and $r \geq 0$. Then*

$$\begin{aligned} & \frac{1}{2}(M_r(x) + (-1)^a M_r(-x)) - \delta_{r+a,0} = \\ & = \sum_{\rho \in K_{2r+1}} x^{s(\rho)} \sum_{r_1, \dots, r_{f(\rho)} = r - c(\rho)} \left(\sum_{\mathbf{v} \in X_{(-1)^a(\rho)}} 2^{-f(\rho)} z_\rho(\mathbf{v}) \prod_{j=1}^{f(\rho)} (M_{r_j}(x) + (-1)^{v_j} M_{r_j}(-x)) \right). \end{aligned}$$

As a remark, the above theorem yields two equations (for $a = 0$ and $a = 1$) that are linear on $M_r(x)$ and $M_r(-x)$. So, Theorem 4.2 provides a finite algorithm for finding $M_r(x)$ for any given $r \geq 0$, since we have to consider all permutations in \mathfrak{S}_{2r+1} , and to perform certain routine operations with all shapes found so far. Moreover, the amount of search can be decreased substantially due to the following proposition which holds immediately by Theorem 3.4 and Theorem 4.2.

Proposition 4.3. *Let $r \geq 1$, $a \in \{0, 1\}$, and*

$$\rho = 2r - 1, 2r + 1, 2r - 3, 2r, \dots, 2r - 2j - 3, 2r - 2j, \dots, 1, 4, 2.$$

Then the expression

$$x^{s(\rho)} \sum_{r_1, \dots, r_{f(\rho)} = r - c(\rho), r_j \geq 0} \left(\sum_{\mathbf{v} \in X_{(-1)^a(\rho)}} 2^{-f(\rho)} z_\rho(\mathbf{v}) \prod_{j=1}^{f(\rho)} (M_{r_j}(x) + (-1)^{v_j} M_{r_j}(-x)) \right)$$

is given by

$$\sum_{j=a}^{\lfloor (r+2)/2 \rfloor} (-1)^{j-a+1} 2^{-r-2} \binom{r+2}{2j+1-a} x^{2r+1} (M_0(x) - M_0(-x))^j (M_0(x) + M_0(-x))^{r+2-j}.$$

By this proposition, it is sufficient to search only permutations in \mathfrak{S}_{2r} . Besides, using Theorem 4.2 and the case $r = 0$ together with induction on r we get the following result.

Theorem 4.4. *$M_r(x)$ is a rational function on x and $\sqrt{1-4x^2}$ for any $r \geq 0$.*

In view of our explicit results, we have even a stronger conjecture.

Conjecture 4.5. *For any $r \geq 1$, there exist polynomials $A_r(x)$, $B_r(x)$, $C_r(x)$, and $D_r(x)$ with integer coefficients such that*

$$E_r(x) = \frac{1}{4}(A_r(x) + B_r(x)) + \frac{1}{4}C_r(x)(1-4x)^{-r+1/2} + \frac{1}{4}D_r(x)(1-4x^2)^{-r+1/2},$$

and

$$O_r(x) = \frac{1}{4}(A_r(x) - B_r(x)) + \frac{1}{4}C_r(x)(1-4x)^{-r+1/2} - \frac{1}{4}D_r(x)(1-4x^2)^{-r+1/2}.$$

REFERENCES

- [1] N. Alon and E. Friedgut. On the number of permutations avoiding a given pattern. *J. Combin. Theory Ser. A*, 89(1):133–140, 2000.
- [2] M. D. Atkinson. Restricted permutations. *Discrete Math.*, 195(1-3):27–38, 1999.
- [3] M. Bóna. Exact enumeration of 1342-avoiding permutations: a close link with labeled trees and planar maps. *J. Combin. Theory Ser. A*, 80(2):257–272, 1997.
- [4] M. Bóna. The number of permutations with exactly r 132-subsequences is P -recursive in the size! *Adv. in Appl. Math.*, 18(4):510–522, 1997.
- [5] M. Bóna. Permutations avoiding certain patterns: the case of length 4 and some generalizations. *Discrete Math.*, 175(1-3):55–67, 1997.
- [6] M. Bóna. Permutations with one or two 132-subsequences. *Discrete Math.*, 181(1-3):267–274, 1998.
- [7] T. Chow and J. West. Forbidden subsequences and Chebyshev polynomials. *Discrete Math.*, 204(1-3):119–128, 1999.
- [8] M. Fulmek. Enumeration of permutations containing a prescribed number of occurrences of a pattern of length three. *Adv. Appl. Math.*, to appear, math.CO/0112092.
- [9] D.E. Knuth. *The Art of Computer Programming*, 2nd ed. Addison Wesley, Reading, MA (1973).
- [10] T. Mansour. Permutations containing and avoiding certain patterns. In *Formal power series and algebraic combinatorics (Moscow, 2000)*, 704–708. Springer, Berlin, 2000.
- [11] T. Mansour and A. Vainshtein. Restricted 132-avoiding permutations. *Adv. Appl. Math.*, 26:258–269, 2001.
- [12] T. Mansour and A. Vainshtein. Counting occurrences of 132 in a permutation. *Adv. Appl. Math.*, 28(2):185–195, 2002.
- [13] J. Noonan. The number of permutations containing exactly one increasing subsequence of length three. *Discrete Math.*, 152(1-3):307–313, 1996.
- [14] J. Noonan and D. Zeilberger. The enumeration of permutations with a prescribed number of “forbidden” patterns. *Adv. in Appl. Math.*, 17(4):381–407, 1996.
- [15] A. Robertson. Permutations containing and avoiding 123 and 132 patterns. *Discrete Math. Theor. Comput. Sci.*, 3(4):151–154 (electronic), 1999.
- [16] R. Simion and F. W. Schmidt. Restricted permutations. *European J. Combin.*, 6(4):383–406, 1985.
- [17] Z. Stankova. Forbidden subsequences. *Discrete Math.*, 132(1-3):291–316, 1994.
- [18] Z. Stankova. Classification of forbidden subsequences of length 4. *European J. Combin.*, 17(5):501–517, 1996.
- [19] J. West. Generating trees and the Catalan and Schröder numbers. *Discrete Math.*, 146(1-3):247–262, 1995.