

The Genesis of the Macdonald Polynomial Statistics

J. Haglund

Department of Mathematics
University of Pennsylvania
Philadelphia, PA 19104-6395
jhaglund@math.upenn.edu

May 9, 2005

Abstract

Recently M. Haiman, N. Loehr and the author [HHL05b], [HHL05a] proved that for μ a partition of n , the modified Macdonald polynomial $\tilde{H}_\mu[z_1, \dots, z_n; q, t]$ can be expressed as a sum of monomials in the z_i times certain nonnegative integral powers of q, t with direct combinatorial descriptions (i.e. statistics). These powers are generalizations of the classical permutation statistics maj and inv . The result was first conjectured by the author [Hag04a], and was partially motivated by a conjectured formula for the character of the space of diagonal harmonics [HHL⁺05c]. Beyond giving a long-sought after combinatorial formula for Macdonald polynomials, this result has many nice corollaries, including a simple proof of Lascoux and Schützenberger’s formula, involving the statistic cocharge , for Hall-Littlewood polynomials. In this paper we describe the sequence of experimental steps and Maple calculations which led to the discovery of the Macdonald polynomial statistics.

In un recente lavoro in collaborazione con M. Haiman, N. Loehr [HHL05b], [HHL05a] dimostriamo che per una partizione di n , μ , il polinomio di Macdonald modificato $\tilde{H}_\mu[z_1, \dots, z_n; q, t]$ può essere espresso come un polinomio nelle variabili z_i i cui coefficienti sono potenze non negative di q, t . Questo risultato ha una diretta interpretazione combinatoria (o meglio statistica). Gli esponenti dei coefficienti sono una generalizzazione delle classiche statistiche di permutazione maj e inv . L’ipotesi di un tale risultato fu avanzata in [Hag04a] e in parte fu motivata dalla formula, in forma di congettura, per i caratteri dello spazio delle armoniche diagonali [HHL⁺05c]. Questo risultato non solo dà un’interpretazione combinatoria, cercata da lungo tempo, dei polinomi di Macdonald, ma permette di derivare anche altri risultati rilevanti. Per esempio è possibile dare una dimostrazione semplificata della formula di Lascoux and Schützenberger riguardante la statistica cocharge per i polinomi di Hall-Littlewood. In questo articolo descriviamo la sequenza di passaggi sperimentali e di calcoli fatti utilizzando Maple che portano alla scoperta della statistica dei polinomi di Macdonald.

1 Introduction

Given a sequence $\mu = (\mu_1, \mu_2, \dots)$ of nonincreasing, nonnegative integers with $\sum_i \mu_i = n$, we say μ is a partition of n , denoted by either $|\mu| = n$ or $\mu \vdash n$. By adding or subtracting parts of size 0 if necessary, we will always assume partitions of n have exactly n parts. We let $\eta(\mu) = \sum_i (i-1)\mu_i$, and if λ is another partition, set $\tilde{K}_{\lambda,\mu}(q,t) = t^{\eta(\mu)} K_{\lambda,\mu}(q, 1/t)$, where $K_{\lambda,\mu}(q,t)$ is Macdonald's q,t -Kostka polynomial [Mac95, p.354]. We call $\tilde{H}_\mu[Z; q, t] = \sum_{\lambda \vdash |\mu|} s_\lambda \tilde{K}_{\lambda,\mu}(q,t)$ the modified Macdonald polynomial, where $s_\lambda = s_\lambda[Z]$ is the Schur function, the sum is over all $\lambda \vdash |\mu|$, and $Z = z_1, \dots, z_n$. The $\tilde{H}_\mu[Z; q, t]$ can be easily transformed by a plethystic substitution into Macdonald's original symmetric functions $P_\mu[Z; q, t]$. Macdonald defined the P_μ in terms of orthogonality with respect to a scalar product, and conjectured $K_{\lambda,\mu}(q,t) \in \mathbb{N}[q, t]$ [Mac95, p. 355]. (From their definition, all one can infer is that the $K_{\lambda,\mu}(q,t)$ are rational functions in q, t). This conjecture in turn led Garsia and Haiman to the “ $n!$ conjecture” [GH93], which was proved by Haiman in 2000 [Hai01]. This result implies that $\tilde{K}_{\lambda,\mu}(q,t) \in \mathbb{N}[q, t]$, and moreover that $\tilde{H}_\mu[X; q, t]$ is the character of a certain S_n -module $V(\mu)$, and thus gives a representation-theoretical interpretation for the coefficients of $\tilde{K}_{\lambda,\mu}(q,t)$. Macdonald also posed the problem of finding a combinatorial rule to describe the $K_{\lambda,\mu}(q,t)$, which is still open.

Recently the author introduced a conjectured combinatorial formula for the coefficient of a monomial in $\tilde{H}_\mu[Z; q, t]$ [Hag04a]. The formula, described in Section 2, is manifestly in $\mathbb{N}[q, t, z_1, \dots, z_n]$. Shortly after its introduction, the conjecture was proved by M. Haiman, N. Loehr and the author. Some comments on the proof, which is surprisingly simple, are made in Section 3. As a corollary they obtain a new, short proof of a famous result of Lascoux and Schützenberger, namely that the coefficients in the Schur function expansion of the Hall-Littlewood polynomial $\tilde{H}_\mu[Z; 0, t]$ can be expressed as a sum, over semi-standard young tableaux W , of q to a statistic known as *cocharge*(W). This new proof is discussed in more detail in Section 4. Also, since the matrix expressing the Schur basis in terms of the monomial basis is upper unitriangular, another consequence is a new proof that the $\tilde{K}_{\lambda,\mu}(q,t) \in \mathbb{Z}[q, t]$. Another consequence is a new proof of Knop and Sahi's formula for Jack symmetric functions [KS97], as well as an extension of their formula to Macdonald's integral form $J_\mu[Z; q, t]$. In addition a simple formula for the $\tilde{K}_{\lambda,\mu}(q,t)$ is obtained when μ has two columns. It is hoped that further refinements will result in combinatorial formulas for the $\tilde{K}_{\lambda,\mu}(q,t)$ for general μ .

The idea for this article was suggested by A. Garsia, who thought it would be beneficial to have a detailed description of the sequence of steps which led the author to the discovery of the Macdonald polynomial statistics. In Section 5 we overview the pioneering work of Garsia, Haiman and others on the $n!$ conjecture, the space of diagonal harmonics DH_n , and the q,t -Catalan numbers. These numbers were defined by Garsia and Haiman as a complicated sum of rational functions in q, t , which they conjectured simplified to a polynomial in $\mathbb{N}[q, t]$ [GH96]. We then discuss a series of discoveries by the author, M. Haiman and others involving combinatorial formulas for the q,t -Catalan numbers and other sequences connected to DH_n , which culminated in the introduction of the “shuffle conjecture” by Haiman, Haglund, Loehr, Remmel and Ulyanov [HHL⁺05c], which gives a combinatorial formula for the character of DH_n .

In Section 6 we discuss how various special cases of the shuffle conjecture led the author to suspect an analogous combinatorial formula for the monomial expansion of $\tilde{H}_\mu[X; q, t]$ could be found. We then include an explanation of the various stages and Maple calculations in the experimental process which resulted in the discovery of the statistics.

2 The Formula

We assign (row,column)-coordinates to squares in the first quadrant, obtained by permuting the (x,y) coordinates of the upper right-hand corner of the square, so the lower left-hand square has coordinates $(1,1)$, the square above it $(2,1)$, etc.. For a square w , we call the first coordinate of w the *row value* of w , denoted $\text{row}(w)$, and the second coordinate of w the *column value* of w , denoted $\text{col}(w)$. Given $\mu \vdash n$, we let μ also stand for the Ferrers diagram of μ (French convention), consisting of the set of n squares with coordinates (i,j) , with $1 \leq i \leq n, 1 \leq j \leq \mu_i$.

Let T be a finite set of squares in the first quadrant. A subset of squares of T consisting of all those $w \in T$ with a given row value is called a row of T , and a subset of squares of T consisting of all those $w \in T$ with a given column value is called a column of T . Furthermore, we let $T(i)$ denote the i th square of T encountered if we read across rows from left to right, starting with the squares of largest row value and working downwards. Given a square $w \in T$, define the leg of w , denoted $\text{leg}(w)$, to be the number of squares in T which are strictly above and in the same column as w , and the arm of w , denoted $\text{arm}(w)$, to be the number of squares in T strictly to the right and in the same row as w . Also, if w has coordinates (i, j) , we let $\text{south}(w)$ denote the square with coordinates $(i - 1, j)$.

A word σ of length n in an alphabet \mathcal{A} is a linear sequence $\sigma_1\sigma_2\cdots\sigma_n$, with $\sigma_i \in \mathcal{A}$. Note that repeats are allowed. If the letter i occurs α_i times in σ , for each $i \geq 1$, we say σ has content α , denoted $\text{content}(\sigma) = \alpha$. We call a pair (σ, T) , where σ is a word of positive integers and T is a set of squares in the first quadrant, a *filling*. We represent (σ, T) geometrically by placing σ_i in square $T(i)$, for $1 \leq i \leq n$. For $w \in T$, we let $\sigma(w)$ denote the element of σ placed in square w , and we call $\sigma_1 \cdots \sigma_n$ the *reading word* of (σ, T) . A *descent* of (σ, T) is a square $w \in T$, with $\text{south}(w) \in T$ and $\sigma(w) > \sigma(\text{south}(w))$.

Let $\text{Des}(\sigma, T)$ denote the set of all descents of (σ, T) . For partitions μ , define a generalized major index statistic $\text{maj}(\sigma, \mu)$ via

$$\text{maj}(\sigma, \mu) = \sum_{w \in \text{Des}(\sigma, \mu)} 1 + \text{leg}(w). \tag{1}$$

An inversion of (σ, T) is a pair of squares (a, b) with $a, b \in T$, $a(\sigma) > b(\sigma)$, and either

$$\begin{cases} \text{row}(a) = \text{row}(b) \text{ and } \text{col}(a) < \text{col}(b), & \text{or} \\ \text{row}(a) = \text{row}(b) + 1 \text{ and } \text{col}(a) > \text{col}(b). \end{cases} \tag{2}$$

Let $\text{Inv}(\sigma, T)$ denote the set of all inversions of (σ, T) , and define the inversion statistic $\text{inv}(\sigma, T)$ via

$$\text{inv}(\sigma, T) = |\text{Inv}(\sigma, T)| - \sum_{w \in \text{Des}(\sigma, T)} \text{arm}(w), \tag{3}$$

where $|T|$ denotes the cardinality of a set T . For example, if (σ, μ) is the filling in Figure 1, then representing squares by their coordinates,

$$\text{Des}(\sigma, \mu) = \{(2, 1), (4, 2), (2, 2)\}, \tag{4}$$

$$\text{Inv}(\sigma, \mu) = \{((3, 1), (3, 2)), ((2, 2), (2, 3)), ((2, 2), (1, 1)), ((2, 3), (1, 1)), ((2, 3), (1, 2))\}, \tag{5}$$

so $\text{maj}(\sigma, \mu) = 3 + 1 + 3 = 7$, $\text{inv}(\sigma, \mu) = 5 - (2 + 0 + 1) = 2$.

2	2	
2	1	
3	5	3
1	1	4

Figure 1: A filling of the partition $(3, 3, 2, 2)$ by the word 2221353114.

Note that, if 1^n denotes a column of n cells, then

$$\text{maj}(\sigma, 1^n) = \sum_{i \in \text{Des}(\sigma, 1^n)} i, \tag{6}$$

the usual major index statistic on the word σ , while

$$\text{inv}(\sigma, (n)) = \sum_{\substack{1 \leq i < j \leq n \\ \sigma_i > \sigma_j}} 1, \tag{7}$$

the usual inversion statistic.

For $\mu \vdash n$, define

$$\tilde{C}_\mu[Z; q, t] = \sum_{\sigma} t^{\text{maj}(\sigma, \mu)} q^{\text{inv}(\sigma, \mu)} z^\sigma, \tag{8}$$

where $z^\sigma = \prod_{i=1}^n z_{\sigma_i}$ is the “weight” of σ and the sum is over all words σ of n positive integers satisfying $1 \leq \sigma_i \leq n$ for $1 \leq i \leq n$. The following result was conjectured by the author [Hag04a] and proven by Haglund, Haiman, Loehr [HHL05b], [HHL05a]).

Theorem 1 *For all partitions μ ,*

$$\tilde{C}_\mu[Z; q, t] = \tilde{H}_\mu[Z; q, t]. \tag{9}$$

Given a set T of squares and a subset $S \subseteq T$, define

$$F_T[Z; q, S] = \sum_{\substack{\sigma \\ \text{Des}(\sigma, T) = S}} q^{\text{inv}(\sigma, T)} z^\sigma. \tag{10}$$

In [Hag04a] the following result is obtained.

Theorem 2 *For all S, T , $F_T[Z; q, S]$ is a symmetric function in the z_i .*

Given $S \subseteq \mu$, let

$$P(S) = \sum_{w \in S} 1 + \text{leg}(w). \tag{11}$$

Note that by the definition of $\text{maj}(\sigma, \mu)$,

$$\tilde{C}_\mu[Z; q, t] = \sum_{S \subseteq \mu} t^{P(S)} F_\mu[Z; q, S]. \tag{12}$$

In [HHL05a] it is shown that the $F_T[Z; q, t]$ are special cases of polynomials introduced by Lascoux, Leclerc and Thibon [LLT97], commonly known as LLT polynomials. These are symmetric polynomials whose description involves a tuple of arbitrary skew shapes and whose coefficients depend on q . It has been an open conjecture of theirs that the coefficients of these polynomials when expanded in the Schur basis are in $\mathbb{N}[q]$. Thus we now have the expansion of $\tilde{H}_\mu[Z; q, t]$ into LLT polynomials, and furthermore understanding the positivity of the coefficients of the $\tilde{K}_{\lambda, \mu}(q, t)$ is reduced to understanding the special case of the positivity of LLT polynomials (when each skew-shape in the tuple is a ribbon).

Definition 1 *Given a word σ of content $(\gamma_1, \gamma_2, \dots)$, construct a permutation σ' , the standardization of σ , by replacing the γ_1 1's in σ by the numbers $1, \dots, \gamma_1$, the γ_2 2's in σ by the numbers $\gamma_1 + 1, \dots, \gamma_1 + \gamma_2$, etc., in such a way that, for $i < j$, $\sigma_i \leq \sigma_j$ if and only if $\sigma'_i < \sigma'_j$. For example, if $\sigma = 224123114$ then $\sigma' = 458167239$.*

Remark 1 *At first glance it may seem that $\text{inv}(\sigma, T)$ may not always be nonnegative, but given a square $u \in \text{Des}(\sigma, T)$, for each square v in the same row as u and to the right of u , either $\sigma(u) > \sigma(v)$, or $\sigma(v) > \sigma(\text{south}(u))$, or both. Assume for the moment that σ has distinct entries. If we adopt the convention that for a square $w \notin T$, $\sigma(w) = \infty$, it follows that $\text{inv}(\sigma, T)$ equals the number of triples of squares u, v, w , where $u \in T$, $v \in T$, $\text{row}(u) = \text{row}(v)$, $\text{col}(u) < \text{col}(v)$, and $\text{south}(u) = w$, and if we draw a circle through u, v, w , and read in the σ values of u, v, w in counterclockwise order around the circle, starting at the smallest value, then the three values form a strictly increasing sequence. If σ has repeated entries, first standardize then count triples in (σ', T) as above.*

3 The Proof

Let $p_k(Z) = \sum_i z_i^k$ be the k th power sum. Given a real parameter w , let $p_k[Z(1-w)] = \sum_i z_i^k (1-w^k)$ and $p_k[Z/(1-w)] = \sum_i z_i^k / (1-w^k)$. These are both special cases of plethystic notation, indicated by the square brackets around $Z(1-w)$ and $Z/(1-w)$. For an arbitrary symmetric function $F(Z)$, let $F[Z(1-w)]$ (respectively $F[Z/(1-w)]$) be the result of first expressing $F(Z)$ as a polynomial in the $p_k(Z)$, then replacing each $p_k(Z)$ by $p_k[Z(1-w)]$ (respectively $p_k[Z/(1-w)]$).

The polynomial $\tilde{H}_\mu[Z; q, t]$ can be defined [Hai03] as the unique polynomial satisfying the following axioms, where $\lambda \leq \mu$ refers to the dominance order $\lambda_1 + \dots + \lambda_i \leq \mu_1 + \dots + \mu_i - 1$ for $1 \leq i \leq n$:

$$(M1) \quad \tilde{H}_\mu[Z(q-1)] = \sum_{\lambda \leq \mu'} a_{\lambda, \mu} m_\lambda \text{ for some } a_{\lambda, \mu} \in \mathbb{Q}(q, t)$$

$$(M2) \quad \tilde{H}_\mu[Z(t-1)] = \sum_{\lambda \leq \mu} a_{\lambda, \mu} m_\lambda \text{ for some } a_{\lambda, \mu} \in \mathbb{Q}(q, t)$$

$$(M3) \quad \text{The coefficient of } z_1^n \text{ in the expansion of } \tilde{H}_\mu[Z; q, t] \text{ into monomials equals 1.}$$

We call a filling of μ by arbitrary nonzero integers a “super filling”. We usually represent negative letters, such as $-2, -4, -7$ by “barred letters” $\bar{2}, \bar{4}, \bar{7}$. Assume the letters satisfy the ordering

$$1 < \bar{1} < 2 < \bar{2} < \dots < n < \bar{n}. \tag{13}$$

Given a super filling σ , we define the standardization σ' of σ to be the unique permutation satisfying $\sigma'_i < \sigma'_j$ whenever $\sigma_i < \sigma_j$, or whenever $i < j$ and $\sigma_i = \sigma_j$ with σ_i a positive letter, or whenever $i > j$ and $\sigma_i = \sigma_j$ with σ_i a negative letter. We then extend the definition of maj and inv to super fillings by letting $\text{maj}(\sigma, \mu) = \text{maj}(\sigma', \mu)$ and $\text{inv}(\sigma, \mu) = \text{inv}(\sigma', \mu)$. Furthermore, we let $|\sigma|$ be the filling obtained by replacing each σ_i by its absolute value.

Using some general results about quasi-symmetric functions and the superization of a symmetric function derived in [HHL⁺05c], one easily obtains the fact that

$$\tilde{C}_\mu[Z(q-1); q, t] = \sum_{\sigma} (-1)^{m(\sigma)} q^{p(\sigma) + \text{inv}(\sigma, \mu)} t^{\text{maj}(\sigma, \mu)} z^{|\sigma|} \tag{14}$$

$$\tilde{C}_\mu[Z(t-1); q, t] = \sum_{\sigma} (-1)^{m(\sigma)} q^{\text{inv}(\sigma, \mu)} t^{p(\sigma) + \text{maj}(\sigma, \mu)} z^{|\sigma|} \tag{15}$$

$$\tag{16}$$

where the sum is over all super fillings σ of μ , $m(\sigma)$ is the number of negative letters in σ , and $p(\sigma)$ is the number of positive letters in σ .

We say two squares u, v of a Ferrers shape attack each other if they are either in the same row, or if u is one row above v and in a column strictly to the right of v . Otherwise u, v are said to be nonattacking. We call a super filling (σ, μ) nonattacking if for all $u, v \in \mu$, $|\sigma(u)| = |\sigma(v)|$ implies u, v are nonattacking.

The first step in the proof of Theorem 1 involves the construction of a sign-reversing involution on super fillings of μ which cancels most of the terms in (14). The involution looks for an attacking pair of squares containing 1's or $\bar{1}$'s. If more than one such pair exists, it chooses the last such pair in the reading word, and switches the sign on the first element of the pair in the reading word. One checks that the q, t -weights are preserved. If no attacking pairs containing $1, \bar{1}$'s exist, then you search for attacking pairs containing $2, \bar{2}$'s, etc. The fixed points are super fillings with no attacking pairs, which are easily seen to satisfy the triangularity condition (M1).

So far we have assumed (13) holds, but in fact (14) and (15) hold for *any* fixed total ordering of the alphabet of positive and negative letters. We now construct a second involution assuming the ordering

$$1 < 2 < \dots < n < \bar{n} < \dots < \bar{2} < \bar{1}. \tag{17}$$

Search for the first occurrence of a 1 or $\bar{1}$ in the reading word, ignoring any such letters in the bottom row. If such a $1, \bar{1}$ exists, then switch its sign. If there is no such 1 or $\bar{1}$, look for the first occurrence of a 2 or $\bar{2}$,

ignoring any letters in the bottom two rows, etc. As in the first involution, the q, t weights are preserved. The fixed points are now super fillings with no 1's or $\bar{1}$'s above the bottom row, no 2's or $\bar{2}$'s above the bottom two rows, etc.. Hence the weight $z^{|\sigma|}$ must satisfy $\lambda_1 \leq \mu_1$, $\lambda_1 + \lambda_2 \leq \mu_1 + \mu_2$, etc., and (M2) is satisfied. The proof is completed by noting that if σ is the filling of all 1's, $\text{inv}(\sigma, \mu) = \text{maj}(\sigma, \mu) = 0$, which implies (M3).

4 The Cocharge Formula

In this section we briefly highlight one of the consequences of Theorem 1, which is described in more detail in [HHL05a]. By considering fillings with no inversions, we obtain the following famous result of Lascoux and Schützenberger, where a semi-standard Young tableau of shape λ and weight μ (or $SSYT(\lambda, \mu)$) is a filling of λ by μ_1 -1's, μ_2 -2's, etc., with entries that are weakly increasing across rows and strictly increasing up columns.

Theorem 3

$$\tilde{H}_\mu[Z; 0, t] = \sum_{\lambda} s_{\lambda} \sum_{T \in SSYT(\lambda, \mu)} t^{\text{cocharge}(\text{read}(T))}, \tag{18}$$

where the statistic cocharge is described (in the proof) below.

Proof. We begin by describing the set of fillings with no inversions. Let M_i be an arbitrary multiset of μ_i positive integers, for $1 \leq i \leq \mu'_1$, and consider those fillings of μ where the elements of M_i are placed in the i th row of μ , in any arbitrary order, for each i in the range $1 \leq i \leq \mu'_1$. It turns out there is exactly one of these fillings with no inversions. This filling can be constructed by first filling the bottom row of μ with the elements of M_1 , in nondecreasing order left to right. Then in square $(2, 1)$, place the smallest integer in M_2 which is strictly bigger than $\sigma_{\mathbf{M}}(1, 1)$, if it exists. If all elements of M_2 are less than or equal to $\sigma_{\mathbf{M}}(1, 1)$, then place the smallest element of M_2 in $(2, 1)$. It is easy to see that this will force any triple of squares of the form $\{(2, 1), (2, j), (1, 1)\}$, $j \geq 2$, to form a clockwise circle in the sense of Remark 1. Next remove $\sigma_{\mathbf{M}}(2, 1)$ from M_2 to form M'_2 , and place the smallest element of M'_2 larger than $\sigma_{\mathbf{M}}(1, 2)$ in square $(2, 2)$, if it exists. If not, place the smallest element of M'_2 in square $(2, 2)$. Now iterate the process, moving left to right, in each square of row two placing the smallest remaining element larger than the element in the square just below, if it exists, and otherwise placing the smallest remaining element in the square. The same process is then applied to row 3, comparing elements of M_3 to elements in the second row of $\sigma_{\mathbf{M}}$, then moving on to row 4, etc.. If $\mu = 5531$, $M_1 = \{1, 1, 3, 6, 7\}$, $M_2 = \{1, 2, 4, 4, 5\}$, $M_3 = \{1, 2, 3\}$, and $M_4 = \{2\}$, then $\sigma_{\mathbf{M}}$ is the filling in Figure 2.

2				
3	1	2		
2	4	4	1	5
1	1	3	6	7

Figure 2: A filling with no inversions.

We now construct a word associated to a filling σ we call the cocharge word $\text{cword}(\sigma)$. Initialize $\text{cword}(\sigma)$ to the empty string, then scan through the reading word of σ , starting at the beginning. Whenever a 1 is encountered, adjoin the number of the row containing this 1 to the left of $\text{cword}(\sigma)$. After reaching the end of the reading word, scan through the reading word again, from the beginning, this time looking for 2's. Whenever a 2 is encountered, adjoin the row number to the left of $\text{cword}(\sigma)$ as

before. After finishing the 2's, loop back and scan through the reading word for 3's, etc. For example, if σ is the filling in Figure 2, $\text{cword}(\sigma) = 11222132341123$.

Next we translate the statistic $\text{maj}(\sigma, \mu)$ into a statistic on $\text{cword}(\sigma)$. Note that $\sigma(1, 1)$ corresponds to the rightmost 1 in $\text{cword}(\sigma)$ - denote this 1 by w_{11} . If $\sigma(2, 1) > \sigma(1, 1)$, $\sigma(2, 1)$ corresponds to the rightmost 2 which is left of w_{11} , otherwise it corresponds to the rightmost 2 (in $\text{cword}(\sigma)$). In any case denote this 2 by w_{12} . More generally, the element in $\text{cword}(\sigma)$ corresponding to $\sigma(i, 1)$ is the first i encountered when travelling left from $w_{1,i-1}$, looping around and starting at the beginning of $\text{cword}(\sigma)$ if necessary. To find the subword $w_{21}w_{22} \cdots w_{2\mu'_2}$ corresponding to the second column of σ , we do the same algorithm on the word $\text{cword}(\sigma)'$ obtained by removing the elements $w_{11}w_{12} \cdots w_{1\mu'_1}$ from $\text{cword}(\sigma)$, then remove $w_{21}w_{22} \cdots w_{2\mu'_2}$ and apply the same process to find $w_{31}w_{32} \cdots w_{3\mu'_3}$ etc..

Clearly $\sigma(i, j) \in \text{Des}(\sigma, \mu)$ if and only if w_{ij} occurs to the left of $w_{i,j-1}$ in $\text{cword}(\sigma)$. Thus $\text{maj}(\sigma, \mu)$ is transparently equal to the statistic $\text{cocharge}(\text{cword}(\sigma))$ described in [Man01, pp.48-49]. Well-known properties of the RSK algorithm now imply Theorem 3. \square

5 Diagonal Harmonics and the $n!$ Conjecture

Assume $\mu \vdash n$, w_1, \dots, w_n is an arbitrary ordering of the squares of μ , and set

$$\Delta_\mu(x_1, \dots, x_n, y_1, \dots, y_n) = \left| x_i^{\text{row}(w_j)} y_i^{\text{col}(w_j)} \right|_{1 \leq i, j \leq n}. \tag{19}$$

For example,

$$\Delta_{221} = \begin{vmatrix} 1 & y_1 & x_1 & x_1 y_1 & x_1^2 \\ 1 & y_2 & x_2 & x_2 y_2 & x_2^2 \\ 1 & y_3 & x_3 & x_3 y_3 & x_3^2 \\ 1 & y_4 & x_4 & x_4 y_4 & x_4^2 \\ 1 & y_5 & x_5 & x_5 y_5 & x_5^2 \end{vmatrix} \tag{20}$$

Note that Δ_{1^n} is, up to sign, the Vandermonde determinant in x_1, \dots, x_n .

Next define the Garsia-Haiman module $V(\mu)$ as the linear span over \mathbb{C} of Δ_μ and its partial derivatives of all orders. An element $\sigma = \sigma_1 \cdots \sigma_n$ in the symmetric group S_n acts on a polynomial $f \in V(\mu)$ via the “diagonal action”

$$\sigma f(x_1, \dots, x_n, y_1, \dots, y_n) = f(x_{\sigma_1}, \dots, x_{\sigma_n}, y_{\sigma_1}, \dots, y_{\sigma_n}). \tag{21}$$

Note that $V(\mu) = \bigoplus_{i,j \geq 0} V(\mu)^{(i,j)}$, where $V(\mu)^{(i,j)}$ is the portion of $V(\mu)$ of bihomogeneous (x, y) -degree (i, j) , and that the diagonal action respects this bigrading.

In 2000 Haiman [Hai01] proved the “ $n!$ conjecture”, first posed in [GH93], which says that the dimension of $V(\mu)$ equals $|\mu|!$. It had previously been shown [Hai99] that the $n!$ conjecture implies the coefficient of $q^i t^j$ in $\tilde{K}_{\lambda, \mu}(q, t)$ equals the multiplicity of the irreducible S_n character χ^λ in the character of $V(\mu)^{(i,j)}$ under the diagonal action, or equivalently that $\tilde{H}_\mu[Z; q, t]$ equals the image of the character of $V(\mu)$ under the Frobenius map which sends χ^λ to s_λ . Macdonald’s conjecture that $\tilde{K}_{\lambda, \mu}(q, t) \in \mathbb{N}[q, t]$ follows.

The definition of the statistic $|\text{Inv}(\sigma, \mu)|$ was motivated by the “ dinv ” statistic occurring in recent work on the combinatorics of the space DH_n of diagonal harmonics [HHL⁺05c],[HL]. This space is defined [Hai94] as the linear span over \mathbb{Q} of the set of all polynomials $f(x_1, \dots, x_n, y_1, \dots, y_n)$ satisfying

$$\sum_{i=1}^n \partial^h x_i \partial^k y_i f = 0, \forall h + k > 0. \tag{22}$$

We can write $DH_n = \bigoplus_{i,j \geq 0} DH_n^{(i,j)}$, and the diagonal action (21) respects this bigrading. The modules $V(\mu)$ are S_n -submodules of DH_n .

Let ∇ be the linear operator on symmetric functions defined on the $\tilde{H}_\mu[Z; q, t]$ basis via

$$\nabla \tilde{H}_\mu[Z; q, t] = t^{n(\mu)} q^{n(\mu')} \tilde{H}_{\mu'}[Z; q, t]. \tag{23}$$

Another famous result of Haiman [Hai02] is that the bigraded character of DH_n under the diagonal action is given by ∇e_n , where e_n is the n th elementary symmetric function. This was first conjectured by Garsia and Haiman in the early 1990's, who made a special study of $\langle \nabla e_n, s_{1^n} \rangle$, the coefficient of s_{1^n} in the expansion of ∇e_n into Schur functions. They called $\langle \nabla e_n, s_{1^n} \rangle$ the “ q, t -Catalan sequence”, denoted $C_n(q, t)$, since they were able to prove that $C_n(1, 1) = \binom{2n}{n} / (n + 1)$, the n th Catalan number. As in the case of $\tilde{K}_{\lambda, \mu}(q, t)$, all one can infer from its definition is that $C_n(q, t)$ is a complicated sum of rational functions in q, t .

A Dyck path is a lattice path, consisting of north $(0, 1)$ and east $(1, 0)$ steps, starting at $(0, 0)$ and ending at (n, n) , which never goes below the diagonal $x = y$. The first major step in the path to the Macdonald polynomial statistics was made in 2000, when Haglund discovered empirically that $C_n(q, t)$ appeared to be expressible as the sum of $q^{\text{area}} t^{\text{bounce}}$ over Dyck paths ifor combinatorial statistics area, bounce [Hag03]. Shortly after, this conjecture was independently discovered by Haiman in the following form, which is more convenient for generalization.

$$C_n(q, t) = \sum_D q^{\text{dinv}(D)} t^{\text{area}(D)}. \tag{24}$$

Here $\text{area}(D)$ is the number of complete squares below D but strictly above $y = x$. We let $a_i = a_i(D)$ denote the number of these squares in the i th row, from the bottom of the square $n \times n$ grid, and define $\text{dinv}(D)$, the number of “diagonal” inversions of D , as the number of pairs (i, j) with

$$1 \leq i < j \leq n \text{ and } a_i = a_j \text{ or } a_i = a_j + 1. \tag{25}$$

For example, for the path in Figure 3 (ignore the numbers in the grid for the moment), we have $\text{area} = 9, \text{dinv} = 13$.

The Haglund-Haiman conjecture for $C_n(q, t)$ was proven shortly after by Garsia and Haglund [GH01], [GH02], by a complicated application of plethystic symmetric function identities previously developed by Garsia, Bergeron and others through a series of papers [GT96], [BGHT99],[GHT99]. More recently Haglund [Hag04b] used the same techniques to prove a more general result conjectured by Egge, Haglund, Kremer and Killpatrick [EHKK03] giving the coefficient of an arbitrary hook shape in ∇e_n in terms of statistics on lattice paths with diagonal $(1, 1)$ steps also allowed (known as Schröder paths).

A consequence of Haiman's formula for the character of DH_n is the fact that the dimension of DH_n , as a \mathbb{Q} -vector space, is $(n + 1)^{n-1}$. This number is well-known to equal the number of parking functions on n cars, which can be represented geometrically by starting with a Dyck path D , then placing the numbers 1 through n immediately to the right of the North steps of D , with strict decrease down columns. Shortly after the proof of the combinatorial formula for $C_n(q, t)$, Haglund and Loehr [HL] conjectured that the Hilbert series of DH_n equals the sum of $q^{\text{dinv}(P)} t^{\text{area}(P)}$ over all parking functions P on n cars. Here the statistic $\text{area}(P)$ is simply $\text{area}(D)$ for the underlying Dyck path D . If we refer to the number in the j -row as car_j , then $\text{dinv}(P)$ is the number of pairs (i, j) which satisfy the conditions (25), and in addition, if $a_i = a_j$, then $\text{car}_j > \text{car}_i$, while if $a_i = a_j + 1$, then $\text{car}_i > \text{car}_j$. The Haglund-Loehr conjecture is still open.

The research into the combinatorics of DH_n culminated with the discovery of the “shuffle conjecture” about two years ago by Haglund, et. al [HHL⁺05c]. This gives a formula for ∇e_n in terms of statistics on “word parking functions”, which are placements of positive integers in the columns as before, but we allow repeats (but still require strict decrease down columns, and require cars to be no larger than n). A given object P of this type is weighted by $z^P q^{\text{dinv}(P)} t^{\text{area}(P)}$, where $\text{area}(P)$ is again the area of the underlying Dyck path. The description of $\text{dinv}(P)$ involves the *diagonal reading word* $\sigma = \sigma(P)$ of P , which is obtained by reading in the cars along diagonals (lines parallel to $y = x$), outside to in and top to bottom. Next standardize σ as in Remark 1, regard σ' as the diagonal reading word of some P' , then set $\text{dinv}(P) = \text{dinv}(P')$. For example, if P is the word parking function in Figure 3, we have $\sigma = 64641532$,

$\sigma' = 74851632$, $\text{area} = 9$, and $\text{dinv} = 6$, with inversion “pairs” (i, j) of rows equal to $(3, 8)$, $(4, 8)$, $(1, 7)$, $(2, 7)$, $(5, 6)$ and $(3, 4)$ each contributing 1 to dinv . Also, z^P is defined as z^σ .

						4		1
						3		0
			6					2
		4						2
		1						1
6								2
5								1 = a ₂
2								0 = a ₁

Figure 3: A word parking function with the a_i on the right

Conjecture 1 (*The shuffle conjecture - [HHL⁺05c]*)

$$\nabla e_n = \sum_P z^P q^{\text{dinv}(P)} t^{\text{area}(P)}, \tag{26}$$

where the sum is over all word parking functions P with n cars.

An equivalent formulation of the conjecture is that the coefficient of the monomial $z_1^{\lambda_1} z_2^{\lambda_2} \dots$ in ∇e_n equals the sum of $q^{\text{dinv}} t^{\text{area}}$ over all parking functions whose diagonal reading word is a shuffle of increasing sequences of lengths $\lambda_1, \lambda_2, \dots$, hence the phrase “shuffle conjecture”. As is well known, the Hilbert series is the coefficient of $z_1 \dots z_n$ in the expansion of the character into monomials, hence the shuffle conjecture contains Haglund and Loehr’s conjectured formula for the Hilbert series of DH_n as a special case. In [HL] it is shown that it also implies the formula for $C_n(q, t)$, as well as the formula for hook shapes in terms of Schröder paths.

6 Water on Mars

Recall that $\tilde{H}_\mu[Z; q, t]$ is the character of the S_n module $V(\mu)$, which is a submodule of DH_n . Thus we could hope that some version of the shuffle conjecture would hold for $\tilde{H}_\mu[Z; q, t]$. Furthermore, since the statistics in the shuffle conjecture are defined solely in terms of the statistics for the Hilbert series of DH_n , together with the simple operation of standardization, we could hope that discovering statistics for the Hilbert series of $V(\mu)$ would result in a combinatorial formula for the monomial expansion of $\tilde{H}_\mu[Z; q, t]$ through a similar standardization process.

Let $\text{Hilb}_\mu(q, t)$ denote the bigraded Hilbert series of $V(\mu)$. The problem of finding combinatorial statistics for $\text{Hilb}_\mu(q, t)$ was studied by Garsia and Haiman, who obtained statistics when μ has two rows or is a hook shape [GH95]. The author had also made some previous unsuccessful attempts at this problem, but the reasoning in the above paragraph increased the stakes dramatically, and so the author decided to give a more determined attack on the problem.

Since the dimension of $V(\mu)$ is $n!$, it is natural to view the problem in terms of searching for a pair of permutation statistics, which depend on μ , to generate $\text{Hilb}_\mu(q, t)$. Since μ has n squares, permutations are in bijection with fillings of μ by distinct integers, and furthermore from a result of Macdonald [Mac95, p.365, Ex.6] one can easily deduce that

$$\text{Hilb}_\mu(1, t) = \sum_{\sigma \in S_n} t^{\text{maj}(\sigma, \mu)}. \tag{27}$$

We now assume that there exists some q -statistic to match with this maj t -statistic, and try to find it.

For a given Dyck path D , let $t^{\text{area}(D)}F_D(Z; q)$ denote the restriction of the sum on the right-hand-side of (26) to those word parking functions for D . It is proven in [HHL⁺05c] that $F_D(Z; q)$ is a symmetric function (and moreover is a constant power of q times an LLT polynomial, where the elements of the tuple are vertical columns associated with the lengths of the vertical segments of D). Let μ be a partition, and abbreviate μ_1 by ℓ . While working with the F_D , the author proved that if $D = D(\mu)$ is the special type of path consisting of μ'_ℓ vertical steps, followed by μ'_ℓ horizontal steps, followed by $\mu'_{\ell-1}$ vertical steps, followed by $\mu'_{\ell-1}$ horizontal steps, etc., then

$$F_{D(\mu)}(Z; q) = q^{\text{mindinv}(D(\mu))} \sum_{\lambda \vdash n} s_\lambda K_{\lambda', \mu'}(q), \tag{28}$$

where $K_{\lambda, \mu}(q) = q^{n(\mu)} \tilde{K}_{\lambda, \mu}(q^{-1})$ is the charge version of the polynomial from Section 4, and $\text{mindinv}(D)$ equals the minimum value of dinv over all word parking functions for D . This can be proven from recurrences for the $F_{D(\mu)}$, obtained by placing the largest car n at the top of columns, then showing these recurrences are equivalent to those obtained by Garsia and Procesi [GP92] for the Hall-Littlewood polynomials. (An equivalent result is contained in [SSW03], where it is deduced from recurrences of this type for general LLT polynomials).

Since

$$\tilde{H}_\mu[Z; 0, q] = \sum_{\lambda} s_\lambda \tilde{K}_{\lambda, \mu}(q), \tag{29}$$

from the first it seemed that any statistics for the $\tilde{H}_\mu[Z; q, t]$ had to be connected to cocharge in some way. Now by (28) the monomial expansion of

$$\sum_{\lambda \vdash n} s_\lambda K_{\lambda', \mu'}(q) \tag{30}$$

can be obtained by summing $q^{-\text{mindinv}(D(\mu)) + \text{dinv}(W)} z^W$ over all word parking functions W for $D(\mu)$, so one might suspect that a power of q related to dinv is the right thing to insert into (27). A parking function P for $D(\mu)$ can be transformed into a filling (σ, μ) by pushing all squares in the i th column from the left down $i - 1$ squares, removing all empty columns, and finally rotating 180 degrees, as in Figure 4. Furthermore this sends $\text{dinv}(\sigma)$ to $|\text{Inv}(\sigma, \mu)|$. Note that area gets sent to maj, since we have descents everywhere. We are thus led to

$$\sum_{\sigma \in S_n} t^{\text{maj}(\sigma, \mu)} q^{-\text{mindinv}(D(\mu)) + |\text{Inv}(\sigma, \mu)|} \tag{31}$$

as a candidate for $\text{Hilb}_\mu(q, t)$.

Maple computations verified that (31) correctly predicts the coefficient of $t^{n(\mu)}$ (the highest power of t) in $\text{Hilb}_\mu(q, t)$, and that the coefficient of t^0 would be correct if we didn't have the $-\text{mindinv}(D(\mu))$ in the q -power. This led to the hypothesis that $|\text{Inv}(\sigma, \mu)|$ formed an upper bound for the correct q -statistic, and that the key was to find the right thing to subtract from it, something possibly depending on the descent set. The final stage involved a series of Maple calculations, each testing a different idea for something to subtract, combined with various maj-like choices for the t -statistic. Many things worked up through $n = 4$ and for various special cases like $\mu = 2, 1, 1, 1, \dots$, but continually failed for $n = 5$. Here is a sample of the author's daily journal summarizing the experiments:

Journal entry:

On 3/29/04, ran pistolbad.map, which computes inversion in columns for tstat and dinv minus mindinv for qstat. Bombs for $\mu = [2, 2, 1]$. See bomb.out Tried a modification, where comaj of the columns replaced the tstat, same qstat. This also bombed in general, but surprisingly works for $\mu = 221, \mu = 222, \mu = 2221$, and $\mu = 2222$. Bombs for $\mu = 2211$ though.




Figure 4: Transforming a parking function into a filling

At this point the author planned to drop the problem, and return to less speculative projects. One last idea occurred to him though, namely that $\text{mindinv}(D(\mu))$ equals the sum of the arm lengths of the squares not in the bottom row of μ , which led to the idea of subtracting the arm of each descent from $|\text{Inv}(\sigma, \mu)|$. A computer run incorporating this idea then successfully generated $\text{Hilb}_\mu(q, t)$ for all $|\mu| \leq 8$. A subsequent calculations a few days later extended this to $|\mu| \leq 9$, and moreover verified that applying the Hilbert series statistics to the standardization of words generated the entire monomial expansion of $\tilde{H}_\mu[Z; q, t]$. In a phone conversation a few months later, after the conjecture was made public, A. Garsia told the author “You found water on Mars”.

Journal Entry:

On 4/2/04 ran `pistolbbad.map`, which computes inversions in columns for `tstat` and `dinv` minus `mindinv` for `qstat`, where `mindinv` is defined as the sum over all descents, of the number of entries to the right of the top element of the decent pair. Seems to work for μ having two rows, but bombs for $\mu = 2, 2, 1$.

Journal Entry:

Also on 4/2/04, ran `pistolb.map`, which computes `maj` on columns for `tstat`, and uses `mindinv` as described in paragraph just above. Surprisingly, works for all μ with $|\mu| \leq 8$! The run for $n = 9$ took over a week, but finally finished successfully.

Journal Entry:

On 4/6/04 ran program `pistolc.map`, which computes the Macdonald poly from shuffles using the hilbert series `inv` and `maj` stats. Correctly generates the Macdonald poly \tilde{H}_μ for all μ with $|\mu| \leq 8$! From 4/7 thru 4/16 ran `tempc.map`, which also verifies shuffle conjecture for $n=9$.

References

- [BGHT99] F. Bergeron, A. M. Garsia, M. Haiman, and G. Tesler. Identities and positivity conjectures for some remarkable operators in the theory of symmetric functions. *Methods Appl. Anal.*, 6:363–420, 1999.
- [EHKK03] E. Egge, J. Haglund, K. Killpatrick, and D. Kremer. A Schroder generalization of Haglund’s statistic on Catalan paths. *Electron. J. Combin.*, 10:Research Paper 16, 21 pp. (electronic), 2003.
- [GH93] A. M. Garsia and M. Haiman. A graded representation model for Macdonald polynomials. *Proc. Nat. Acad. Sci. U.S.A.*, 90(8):3607–3610, 1993.
- [GH95] A. M. Garsia and M. Haiman. Factorizations of Pieri rules for Macdonald polynomials. *Discrete Math.*, 139(1-3):219–256, 1995. Formal power series and algebraic combinatorics (Montreal, PQ, 1992).

- [GH96] A. M. Garsia and M. Haiman. A remarkable q, t -Catalan sequence and q -Lagrange inversion. *J. Algebraic Combin.*, 5(3):191–244, 1996.
- [GH01] A. M. Garsia and J. Haglund. A positivity result in the theory of Macdonald polynomials. *Proc. Nat. Acad. Sci. U.S.A.*, 98:4313–4316, 2001.
- [GH02] A. M. Garsia and J. Haglund. A proof of the q, t -Catalan positivity conjecture. *Discrete Math.*, 256:677–717, 2002.
- [GHT99] A. M. Garsia, M. Haiman, and G. Tesler. Explicit plethystic formulas for Macdonald q, t -Kostka coefficients. *Sém. Lothar. Combin.*, 42:Art. B42m, 45 pp. (electronic), 1999. The Andrews Festschrift (Maratea, 1998).
- [GP92] A. M. Garsia and C. Procesi. On certain graded S_n -modules and the q -Kostka polynomials. *Adv. Math.*, 94(1):82–138, 1992.
- [GT96] A. M. Garsia and G. Tesler. Plethystic formulas for Macdonald q, t -Kostka coefficients. *Adv. Math.*, 123:144–222, 1996.
- [Hag03] J. Haglund. Conjectured statistics for the q, t -Catalan numbers. *Adv. Math.*, 175(2):319–334, 2003.
- [Hag04a] J. Haglund. A combinatorial model for the Macdonald polynomials. *Proc. Nat. Acad. Sci. U.S.A.*, 101:16127–16131, 2004.
- [Hag04b] J. Haglund. A proof of the q, t -Schroder conjecture. *Internat. Math. Res. Notices*, 11:525–560, 2004.
- [Hai94] M. Haiman. Conjectures on the quotient ring by diagonal invariants. *J. Algebraic Combin.*, 3:17–76, 1994.
- [Hai99] M. Haiman. Macdonald polynomials and geometry. In *New perspectives in algebraic combinatorics (Berkeley, CA, 1996–97)*, volume 38 of *Math. Sci. Res. Inst. Publ.*, pages 207–254. Cambridge Univ. Press, Cambridge, 1999.
- [Hai01] M. Haiman. Hilbert schemes, polygraphs, and the Macdonald positivity conjecture. *J. Amer. Math. Soc.*, 14:941–1006, 2001.
- [Hai02] M. Haiman. Vanishing theorems and character formulas for the Hilbert scheme of points in the plane. *Invent. Math.*, 149:371–407, 2002.
- [Hai03] Mark Haiman. Combinatorics, symmetric functions, and Hilbert schemes. In *Current developments in mathematics, 2002*, pages 39–111. Int. Press, Somerville, MA, 2003.
- [HHL05a] J. Haglund, M. Haiman, and N. Loehr. A combinatorial formula for Macdonald polynomials. *Jour. Amer. Math. Soc.*, to appear, 2005.
- [HHL05b] J. Haglund, M. Haiman, and N. Loehr. Combinatorial theory of Macdonald polynomials I: Proof of Haglund’s formula. *Proc. Nat. Acad. Sci. U.S.A.*, 102(8):2690–2696, 2005.
- [HHL⁺05c] J. Haglund, M. Haiman, N. Loehr, J. B. Remmel, and A. Ulyanov. A combinatorial formula for the character of the diagonal coinvariants. *Duke J. Math.*, 126:195–232, 2005.
- [HL] J. Haglund and N. Loehr. A conjectured combinatorial formula for the Hilbert series for Diagonal Harmonics. *Discrete Math.*, to appear. Proceedings of the FPSAC 2002 Conference, preprint available at www.math.upenn.edu/~jhaglund/.
- [KS97] Friedrich Knop and Siddhartha Sahi. A recursion and a combinatorial formula for Jack polynomials. *Invent. Math.*, 128(1):9–22, 1997.

- [LLT97] Alain Lascoux, Bernard Leclerc, and Jean-Yves Thibon. Ribbon tableaux, Hall-Littlewood functions, quantum affine algebras, and unipotent varieties. *J. Math. Phys.*, 38(2):1041–1068, 1997.
- [Mac95] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford Mathematical Monographs, second ed., Oxford Science Publications. The Clarendon Press Oxford University Press, New York, 1995.
- [Man01] Laurent Manivel. *Symmetric functions, Schubert polynomials and degeneracy loci*, volume 6 of *SMF/AMS Texts and Monographs*. American Mathematical Society, Providence, RI, 2001. Translated from the 1998 French original by John R. Swallow, Cours Spécialisés [Specialized Courses], 3.
- [SSW03] A. Schilling, M. Shimozono, and D. E. White. Branching formula for q -Littlewood-Richardson coefficients. *Adv. in Appl. Math.*, 30(1-2):258–272, 2003. Formal power series and algebraic combinatorics (Scottsdale, AZ, 2001).

