q-WZ-Theory and Bailey Chains

Peter Paule
RISC, J. Kepler University of Linz, Austria

February 16, 1998

[summary by Bruno Gauthier and Bruno Salvy]

Abstract

Many combinatorial identities can be formulated in terms of q-hypergeometric sums, for instance, the celebrated Rogers-Ramanujan identities from additive number theory. Identities of this type can be constructed iteratively from simpler ones, i.e., by proceeding along Bailey chains. Another construction mechanism, different from this classical one, arises within the context of q-WZ-theory. For instance, as a by-product of computer proofs, one automatically obtains the so-called "dual" identities. The talk gives a short tutorial introduction and discusses various relations between these concepts.

The talk consists of four parts. The first part is an introduction to Gaussian polynomials. The second part is a brief account of q-hypergeometric WZ theory. The parts that follow are variations on this theme.

1. Gaussian polynomials

Let $p(m,n,k)$ denote the number of permutations of k in at most m parts, each part $\leq n$. Clearly,

$$p(m,n,k) = 0, \quad \text{if } k > mn,$$
$$p(m,n,mn) = 1.$$

Therefore the generating function $G_{m,n}(q) = \sum_{k=0}^{mn} p(m,n;k)q^k$ is a polynomial in q of degree mn.

A few particular instances are:

$$G_{m,0}(q) = 1, \qquad G_{0,n}(q) = 1, \qquad G_{m,n}(q) = G_{n,m}(q), \qquad G_{m,1}(q) = \frac{1-q^{m+1}}{1-q},$$

$$G_{4,3}(q) = 1 + q + 2q^2 + 3q^3 + 4q^4 + 4q^5 + 5q^6 + 4q^7 + 4q^8 + 3q^9 + 2q^{10} + q^{11} + q^{12}.$$

From the decomposition

$$p(m,n;k) = p(m-1,n;k-n) + p(m,n-1;k) \qquad (k \geq n)$$

follows that

$$G_{m,n}(q) = q^n G_{m-1,n}(q) + G_{m,n-1}(q). \tag{1}$$

By symmetry, we also have:

$$G_{m,n}(q) = G_{m-1,n}(q) + q^m G_{m,n-1}(q). \tag{2}$$

So, by elimination between (1) and (2):

$$G_{m,n}(q) = \frac{1-q^{m+n}}{1-q^n} G_{m,m-1}(q) = \frac{(1-q^{m+n})\cdots(1-q^{m+1})}{(1-q^n)\cdots(1-q)} G_{m,0}(q).$$

Using the standard notation $(a;q)_k = (1-a)(1-aq)\cdots(1-aq^{k-1})$, $k = 1, 2, \ldots$ and $(a;q)_k = 1/(a;q)_{-k}$ for $k < 0$, we get a closed form representation of the Gaussian polynomials $G_{m,n}$:

$$\begin{bmatrix} m+n \\ n \end{bmatrix} := G_{m,n}(q) = \frac{(1-q^{m+n})\cdots(1-q^{m+1})}{(1-q^n)\cdots(1-q)} = \frac{(q;q)_{m+n}}{(q;q)_m (q;q)_n}. \tag{3}$$

2. Some facts about q-hypergeometric WZ theory

Definition 1. A sequence (t_k) is *hypergeometric* if the ratio of two consecutive terms is a rational function of the summation index k: $t_{k+1}/t_k = P(k)/Q(k)$, where P and Q are polynomials in k.

Definition 2. A sequence (t_k) is *q-hypergeometric* if the ratio of two consecutive terms is a rational function of q^k: $t_{k+1}/t_k = P(q^k)/Q(q^k)$, where P and Q are polynomials in q^k. (Note that q should be contained in the coefficient field which should be of characteristic 0.)

2.1. **From Gosper to Zeilberger.** Gosper's algorithm for indefinite hypergeometric summation [3] is given as input a hypergeometric sequence (f_k). This algorithm finds a hypergeometric sequence (g_k) such that $f_k = g_{k+1} - g_k$ (then g_k is the product of f_k and a rational function in k). From there by telescoping one gets:

$$\sum_{k=0}^{n} f_k = g_{n+1} - g_0.$$

Zeilberger's algorithm computes definite hypergeometric sums. Given a proper hypergeometric sequence $(F_{n,k})$ (with finite support in k), this algorithm finds a hypergeometric sequence (S_k) such that

$$\sum_k F_{n,k} = S_n.$$

The idea is use an extension of Gosper's algorithm in order to find polynomials $a_j(n)$ and a proper hypergeometric term $(G_{n,k})$ such that

$$\sum_{j=0}^{J} a_j(n) F_{n+j,k} = G_{n,k+1} - G_{n,k}.$$

Then $G_{n,k}$ is necessarily of the form $R(n,k)F_{n,k}$ where R is a rational function called the "certificate". Summing this equality yields the desired recurrence on S_n:

$$\sum_{j=0}^{J} a_j(n) S_{n+j} = 0.$$

This algorithm has been implemented in Mathematica by P. Paule and M. Schorn:

Example.

```
In[1]:= <<zb_alg.m
Fast Zeilberger by Peter Paule and Markus Schorn. (V 2.2)
Systembreaker = ENullspace
In[2]:= Zb[Binomial[n,k] x^k, k, n, 1]
Out[2]= {(1 + x) F[k, n] - F[k, 1 + n] == Delta[k, R F[k, n]]}
In[3]:= Show[R]

            k
Out[3]= ---------
        1 - k + n
```

Both these algorithms extend to the q-case, a corresponding implementation in Mathematica is due to A. Riese.

2.2. **Example.** A variation of a q-analogue of Gosper's and Zeilberger's algorithms can serve in finding q-analogues of binomial identities. For instance, in order to derive a q-analogue of the binomial theorem, the question is to find $\alpha, \beta \in \mathbb{Z}$ such that the following sequence satisfies a recurrence of order one:

$$S_n(q) = \sum_{k=0}^{n} \begin{bmatrix} n \\ k \end{bmatrix} x^k q^{\alpha\binom{k}{2}+\beta k},$$

Riese's Mathematica package `qZeil` automatically determines the candidates $\alpha \in \{1\}$, $\beta \in \mathbb{Z}$. Indeed, choosing $\alpha = 1$ and $\beta = 0$:

```
In[3]:= <<qZeil.m

Out[3]= Axel Riese's q-Zeilberger implementation version 1.8 loaded

In[4]:= qZeil[ qBinomial[n,k,q] x^k q^Binomial[k,2], {k,0,n}, n, 1]
                              -1 + n
Out[4]= SUM[n] == -((-1 - q       x) SUM[-1 + n])
```

So, we obtain the q-binomial theorem in one of its standard forms:

$$\sum_{k=0}^{n} \begin{bmatrix} n \\ k \end{bmatrix} x^k q^{\binom{k}{2}} = (1+x)(1+qx)\cdots(1+q^{n-1}x).$$

2.3. **WZ duality.** Given a hypergeometric sequence $(f_{n,k})$, assume that Zeilberger's algorithm finds a_1, a_2 and $g_{n,k}$ such that:

$$a_1(n) f_{n+1,k} + a_2(n) f_{n,k} = g_{n,k+1} - g_{n,k}.$$

Then, in case of finite support:

$$a_1(n) S_{n+1} + a_2(n) S_n = 0.$$

By using this in the form $a_2(n)/S_{n+1} = -a_1(n)/S_n$, we rewrite the relation above as:

$$\frac{a_1(n)}{S_n}\frac{f_{n+1,k}}{S_{n+1}} + \frac{a_2(n)}{S_{n+1}}\frac{f_{n,k}}{S_n} = \frac{g_{n,k+1} - g_{n,k}}{S_{n+1}S_n}.$$

Defining $F_{n,k} = f_{n,k}/S_n$, and $G_{n,k} = g_{n,k}/(a_1(n)S_{n+1})$, we arrive at:

$$F_{n+1,k} - F_{n,k} = G_{n,k+1} - G_{n,k}. \tag{4}$$

This gives rise to the following definition:

Definition 3. A pair of sequences (F, G) that satisfy (4) is called a "WZ pair".

Note that given such a WZ-pair (F having finite support), from (4) follows that $\sum_k F_{n+1,k} - \sum_k F_{n,k} = 0$, which means that the corresponding sum sequence $S_n := \sum_k F(n,k)$ is a constant.

3. A Fibonacci q-analogue

The well-known Fibonacci numbers are defined by $F_0 = 1, F_1 = 1$ and $F_{n+2} = F_{n+1} + F_n$. Does there exist a q-analogue of these numbers? In order to follow the strategy explained above (example 2.2), we take as a starting point the following well-known hypergeometric sum:

$$F_n = \sum_{k=0}^{n} \binom{n-k}{k}.$$

For the (α,β)-Ansatz, we take:

$$F_n(q)=\sum_{k=0}^{n}\begin{bmatrix}n-k\\ k\end{bmatrix}q^{\alpha\binom{k}{2}+\beta k},$$

and we want to determine $\alpha,\beta\in\mathbb{Z}$ such that $(F_n(q))$ satisfies a linear recurrence of order 2. Riese's implementation delivers as candidates: $\alpha\in\{1,2,3\}$ and $\beta\in\mathbb{Z}$, but only the choice $\alpha=2$ is successful. This means, only for $\alpha=2$, the q-analogue of Zeilberger's algorithm delivers ($\forall\beta\in\mathbb{Z}$) a recurrence of order 2, namely:

$$F_{n+2}(q)=F_{n+1}(q)+q^{n+\beta}F_n(q). \tag{5}$$

Let us fix (for instance) $\beta=1$, and we obtain for this choice:

$$F_n(q)=\sum_{k=0}^{n}\begin{bmatrix}n-k\\ k\end{bmatrix}q^{k^2}.$$

In the limit $n\to\infty$:

$$F_\infty(q)=\sum_{k=0}^{\infty}\frac{q^{k^2}}{(q;q)_k}=1+\sum_{n=1}^{\infty}b_nq^n=\prod_{k=1}^{\infty}\frac{1}{(1-q^{5n+1})(1-q^{5n+4})}, \tag{6}$$

where b_n is the number of partitions of n into parts with minimal difference two and the right-hand side is one of the celebrated Rogers-Ramanujan identities [1].

Starting from (5), it is also possible to conjecture and then prove (by q-Zeilberger) the following identity due to I. Schur

$$F_{2n}(q)=\sum_k q^{k(10k+1)}\begin{bmatrix}2n\\ n-5k\end{bmatrix}-\sum_k q^{(2k-1)(5k-2)}\begin{bmatrix}2n\\ n-5k+2\end{bmatrix}.$$

4. **The Bailey chain approach**

Proposition 1.

$$\sum_{j=0}^{n}\frac{q^{j^2-k^2}}{(q;q)_{n-j}}\frac{(q;q)_{n-k}(q;q)_{n+k}}{(q;q)_{j-k}(q;q)_{j+k}}=1. \tag{7}$$

Proof. Denote by $f_{n,j}$ the summand. Riese's implementation yields:

$$f_{n,j}-f_{n-1,j}=g_{n,j}-g_{n,j-1},\qquad\text{where}\qquad g_{n,j}=\frac{q^{k+n}(q^j-q^n)}{(q^n-q^k)(1-q^{k+n})}f_{n,j}.$$

This (q)WZ-pair implies that the sum over $f_{n,j}$ is constant. That this constant is 1 follows from instance from the evaluation for $n=k$ [1, 4]. □

This identity is a special case of a q-hypergeometric formula that can be proved combinatorially as explained in [4].

Multiplying (7) by an arbitrary sequence (c_k), we obtain the following special case of "Bailey's Lemma":

$$\sum_k\frac{c_k}{(q;q)_{n-k}(q;q)_{n+k}}=\sum_{j\ge0}\frac{q^{j^2}}{(q;q)_{n-j}}\sum_k\frac{c_kq^{-k^2}}{(q;q)_{j-k}(q;q)_{j+k}}. \tag{8}$$

Definition 4. Two sequences $((a)_{k\in\mathbb{Z}},(b)_{n\in\mathbb{N}})$ are called a *Bailey-pair* when

$$b_n=\sum_k \frac{a_k}{(q;q)_{n-k}(q;q)_{n+k}}.$$

Now, let's walk in a "Bailey-chain" (using proposition 1) starting with: $a_k=q^{\binom{k}{2}}(-1)^k$ and b_n as above. Using `qZeil`, we get:

$$b_n=\sum_k \frac{q^{\binom{k}{2}}x^k}{(q;q)_{n-k}(q;q)_{n+k}}=\frac{(-x;q)_n(-q/x;q)_n}{(q;q)_{2n}}.$$

Note that in the limit $n\to\infty$, this turns into Jacobi's triple product identity:

$$\sum_k q^{\binom{k}{2}}x^k=\prod_{j=1}^{\infty}(1-q^j)(1+q^{j-1}x)(1+\frac{q^j}{x}).$$

From there (6) follows when substituting q by q^5 and x by $-q^2$.

Now, with $c_k=q^{k^2}a_k$ and $x=-1$, (8) yields an identity due to Rogers:

$$\sum_k \frac{(-1)^k q^{\frac{3}{2}k^2-\frac{1}{2}k}}{(q;q)_{n-k}(q;q)_{n+k}}=\frac{1}{(q;q)_n},$$

which can be found by the q-Zeilberger algorithm after "creative symmetrizing" (i.e., multiplying the summand by $1+q^k$ in this example).

The second step in the Bailey chain approach uses $c_k=q^{2k^2}a_k$. This gives:

$$\sum_k \frac{(-1)^k q^{\frac{5}{2}k^2-\frac{1}{2}k}}{(q;q)_{n-k}(q;q)_{n+k}}=\sum_{j\geq 0}\frac{q^{j^2}}{(q;q)_{n-j}(q;q)_j}.$$

In the limit $n\to\infty$ and by Jacobi's triple product identity, this gives again (6). The third step of the Bailey chain gives:

$$\sum_k \frac{(-1)^k q^{\frac{7}{2}k^2-\frac{1}{2}k}}{(q;q)_{n-k}(q;q)_{n+k}}=\sum_{j\geq 0}\frac{q^{j^2}}{(q;q)_{n-j}}\sum_{l=0}^{j}\frac{q^{l^2}}{(q;q)_{j-l}(q;q)_l},$$

resulting when $n\to\infty$ in an identity due to B. Gordon

$$\frac{1}{(q;q)_\infty}\prod_{n=0}^{\infty}(1-q^{7n+3})(1-q^{7n+4})(1-q^{7n+7})=\sum_{l,j=0}^{\infty}\frac{q^{(j+l)^2+l^2}}{(q;q)_j(q;q)_l},$$

whose first automatic proof was given by Chyzak [2].

References

[1] Andrews (George E.). – q-series : Thier development and application in analysis. *CBMS Regional Conference Series*, n° 66, 1986.

[2] Chyzak (Frédéric). – Groebner bases, symbolic summation and symbolic integration. In Buchberger (B.) and Winkler (F.) (editors), *Groebner Bases and Applications (Proc. of the Conference 33 Years of Gröbner Bases)*. – Cambridge University Press, 1998. ISBN 0 521 63298 6.

[3] Gosper (R. William). – Decision procedure for indefinite hypergeometric summation. *Proceedings of the National Academy of Sciences USA*, vol. 75, n° 1, January 1978, pp. 40–42.

[4] Paule (Peter). – Ueber das inversionsprinzip von garsiaund milne. *Bayreuther Mathematische Schriften*, vol. 21, 1986, pp. 295–319.

[5] Paule (Peter) and Riese (Axel). – A Mathematica q-analogue of Zeilberger's algorithm based on an algebraically motivated approach to q-hypergeometric telescoping. In *Special Functions, q-Series and Related Topics, Toronto, Ontario, 1995. Fields Inst. Commun.*, vol. 14, pp. 179–210. – Providence, 1997.

[6] Paule (Peter) and Schorn (Markus). – A Mathematica version of Zeilberger's algorithm for proving binomial coefficient identities. *Journal of Symbolic Computation*, vol. 20, 1995, pp. 673–698.

[7] Petkovšek (Marko), Wilf (Herbert), and Zeilberger (Doron). – *A=B*. – A. K. Peters, Ltd., Wellesley, Massachusset, 1996. ISBN 1-56881-063-6.